

Coevolution of metal availability and nitrogen assimilation in cyanobacteria and algae

J. B. GLASS,¹ F. WOLFE-SIMON^{2,3} AND A. D. ANBAR^{1,3}

¹*School of Earth and Space Exploration, Arizona State University, Tempe AZ 85287, USA*

²*Department of Earth and Planetary Sciences, Harvard University, Cambridge, MA 02467, USA, and*

³*Department of Chemistry and Biochemistry, Arizona State University, Tempe, AZ 85287, USA*

ABSTRACT

Marine primary producers adapted over eons to the changing chemistry of the oceans. Because a number of metalloenzymes are necessary for N assimilation, changes in the availability of transition metals posed a particular challenge to the supply of this critical nutrient that regulates marine biomass and productivity. Integrating recently developed geochemical, biochemical, and genetic evidence, we infer that the use of metals in N assimilation – particularly Fe and Mo – can be understood in terms of the history of metal availability through time. Anoxic, Fe-rich Archean oceans were conducive to the evolution of Fe-using enzymes that assimilate abiogenic NH_4^+ and NO_2^- . The N demands of an expanding biosphere were satisfied by the evolution of biological N_2 fixation, possibly utilizing only Fe. Trace O_2 in late Archean environments, and the eventual 'Great Oxidation Event' c. 2.3 Ga, mobilized metals such as Mo, enabling the evolution of Mo (or V)-based N_2 fixation and the Mo-dependent enzymes for NO_3^- assimilation and denitrification by prokaryotes. However, the subsequent onset of deep-sea euxinia, an increasingly-accepted idea, may have kept ocean Mo inventories low and depressed Fe, limiting the rate of N_2 fixation and the supply of fixed N. Eukaryotic ecosystems may have been particularly disadvantaged by N scarcity and the high Mo requirement of eukaryotic NO_3^- assimilation. Thorough ocean oxygenation in the Neoproterozoic led to Mo-rich oceans, possibly contributing to the proliferation of eukaryotes and thus the Cambrian explosion of metazoan life. These ideas can be tested by more intensive study of the metal requirements in N assimilation and the biological strategies for metal uptake, regulation, and storage.

Received 21 September 2008; accepted 11 January 2009

Corresponding author: Jennifer Glass. Tel.: 480-244-3603; fax: 262-244-3605; e-mail: jennifer.b.glass@asu.edu

INTRODUCTION

All living things require N. This element is an essential constituent of amino acids, nucleic acids, pigments such as chlorophyll, and other biomolecules. Although N is the most abundant element in the atmosphere, it is overwhelmingly present as dinitrogen (N_2), which is not particularly soluble. N is therefore relatively scarce in the oceans, with a total concentration of <600 micromolar (μM) (Sharp, 1983). The acquisition of N is therefore a major ecological challenge for marine biota.

The dominant primary producers in the modern marine environment are photoautotrophs. Such organisms have probably dominated since the evolution of photosynthesis at least 2.7

billion years ago (Brocks *et al.*, 1999). Today, photoautotrophs include both prokaryotes (cyanobacteria) and eukaryotes (algae). Both types of organisms have evolved a variety of N assimilation pathways to take advantage of the variety of N species in the environment. Today, these N species include:

- Dissolved N_2 , the dominant form of N in the oceans, other than particulate N bound up in living and dead organisms. However, it is also highly nonreactive because of the strength of the N-N bond. Breaking this triple bond is energetically expensive, and so the organisms that can assimilate N_2 only do so when there are no other options.
- Dissolved ammonium (NH_4^+), which is the most biologically accessible form of inorganic N because it is already in the chemical form useful for biomolecules; no energetically expensive reduction reactions are required for its assimilation. However, the concentration of NH_4^+ is exceedingly low in

For Special Issue of Geobiology, 'Co-evolution of Plants and the Environment,' edited by D. Beerling

present-day oxygenated oceans, typically $<0.1 \mu\text{M}$ (Sharp, 1983), as it is rapidly converted to other forms of N by biologically mediated oxidation and other processes.

- Oxidized N species, particularly nitrate (NO_3^-), are more abundant than NH_4^+ . Collectively, these dissolved inorganic forms of N are known as DIN. Although the concentration of NO_3^- is low in surface waters ($0.2 \mu\text{M}$) as a result of biological removal, it can reach up to $\sim 40 \mu\text{M}$ in deep waters (Sharp, 1983). Thus, upwelling events contribute major influxes of NO_3^- to the photic zone. A nitrite (NO_2^-) maximum exists at the base of the euphotic zone in stratified water columns, where the NO_2^- concentration can reach $4.5 \mu\text{M}$ (Lomas & Lipschultz, 2006).
- Dissolved organic nitrogen (DON), excreted by bacteria and zooplankton (Bronk *et al.*, 2007), is the other major form of N. Many nitrogenous compounds compose the DON pool. Urea and amino acids are among the most common constituents of this pool. In the open ocean, these compounds are present at concentrations of $\sim 0.1\text{--}0.5$ and $0.01\text{--}0.03 \mu\text{M}$, respectively (Antia *et al.*, 1991). Urea runoff from agricultural lands has led to elevated urea concentrations in coastal regions, where it contributes to eutrophication (Glibert *et al.*, 2006) and is an important N source for toxic algal species (Dyrhman & Anderson, 2003).

Assimilation of these various forms of N requires a number of complex enzymes (Falkowski, 1983; Berges & Mulholland 2008). Many of these enzymes incorporate transition metals such as iron (Fe), molybdenum (Mo), nickel (Ni) and copper (Cu). Cubic clusters of Fe and S constitute a very common type of cofactor in N assimilatory proteins. Chemically similar to pyrite, they occur in Fe : S ratios of 2 : 2, 3 : 4 and 4 : 4, and perform electron transfer and redox reactions. Fe can also be bound at the center of heme in eukaryotic nitrate reductase (as a cytochrome b_{557}) and nitrite reductase (as siroheme). Like Fe-S clusters, heme can perform electron transfer and catalysis. Mo is bound to a large Fe-S cluster in nitrogenase, and is at the heart of molybdopterin cofactors in both prokaryotic and eukaryotic nitrate reductase. Ni and Cu are not bound to organic cofactors, but are present as two- and one-atom centers, respectively, at the active sites of DON assimilatory proteins.

Biogeochemical cycling of N therefore depends strongly on Fe, Mo, Ni, and Cu (Morel *et al.*, 2003; Morel, 2008). The abundances of these elements in the environment are typically far lower than those of N species. Transition metals in the open ocean are generally present in trace concentrations, in the low nanomolar (nM) to picomolar (pM) range, because of their low solubilities and rapid removal on particles (Morel & Price, 2003). Fe, Mo, Ni, and Cu are present in total concentrations of $0.002\text{--}1 \text{ nM}$ (Johnson *et al.*, 1997), 105 nM (Collier, 1985), $2\text{--}12 \text{ nM}$, and $0.5\text{--}4.5 \text{ nM}$ (Bruland, 1980), respectively, in open ocean surface waters. In coastal waters, metal concentrations are typically higher but much more variable. For example, Fe ranges from <1 to $>10 \text{ nM}$ along the California coast (Bruland *et al.*, 2001), coastal Cu concentrations can reach 50 nM (Moffett *et al.*, 1997; Achterberg *et al.*, 1999),

and Mo in the North Sea can fluctuate between 30 and 160 nM (Dellwig *et al.*, 2007). Fe and Cu are generally present at higher concentrations in freshwater than in seawater (Xue *et al.*, 1995; Achterberg *et al.*, 1997), whereas Ni concentrations show little salinity dependence (Xue *et al.*, 2001). Mo exhibits the opposite trend, with very low concentrations ($<1\text{--}20 \text{ nM}$) in freshwater (Sugawara *et al.*, 1961; Bachmann & Goldman, 1964; Howarth & Cole, 1985; Cole *et al.*, 1993).

Because the metals required for N assimilation are scarce, the pathways for N assimilation in the oceans are strongly affected by metal availability. For example, Fe-fertilization experiments reveal that phytoplankton switch to more Fe-intensive N assimilation pathways when Fe is added to ocean regions where Fe scarcity limits primary production (Martin & Fitzwater, 1988; Price *et al.*, 1991; Coale *et al.*, 1996). Marine cyanobacteria have been shown to fix N_2 at higher rates when provided excess Fe (Berman-Frank *et al.*, 2001; Berman-Frank *et al.*, 2007). It has been hypothesized that the availability of other metals may affect N assimilation in the modern oceans. It is possible that Ni scarcity limits urea utilization in the marine environment as a result of organic complexation and slow uptake kinetics (see references in Dupont *et al.*, 2007). It is unlikely that Mo scarcity limits N assimilation in marine environments, because of its long residence time and high stability in oxic seawater, but Mo scarcity may limit N uptake, and hence primary productivity, in freshwater (Goldman, 1960; Axler *et al.*, 1980).

The pathways for N assimilation almost certainly changed over the course of Earth history because the ocean abundances of these bioessential metals are not constant with time. Their abundances have varied in particular as a consequence of secular changes in ocean oxygenation, which affects element delivery to, and residence times in, the oceans. The classic example is Fe. Scarce in the modern oxygenated oceans, geologic evidence indicates that Fe was relatively abundant in the surface and deep oceans during the Archean ($3.8\text{--}2.5$ billions of years ago (Ga)) as a result of widespread ocean anoxia, perhaps reaching concentrations of tens of micromolar (Fig. 1) (Holland, 1973). Strong evidence has emerged recently for comparably large changes in the deep and surface abundance of Mo, but with an opposite sense of direction than Fe (Fig. 1) (Scott *et al.*, 2008). Analogous shifts probably characterized other elements with redox-sensitive environmental chemistries (Saito *et al.*, 2003), although supporting evidence for these less abundant elements is difficult to extract from the geologic record. However, the connections between historical changes in metal availability and the metal requirements of N assimilation, and their evolutionary consequences, have barely been explored (Falkowski, 1997; Anbar & Knoll, 2002; Anbar, 2008).

One example of such a connection can be seen in the marked differences in N assimilation pathways between prokaryotes and eukaryotes. Prokaryotes can fix N_2 , whereas eukaryotes cannot; eukaryotes assimilate fixed forms of N from the environment or (in the case of heterotrophs) from their food.




Fig. 1 Notional estimated surface (dotted lines) and deep (solid lines) concentrations of Fe (red), Mo (blue) and O₂ (black) in seawater through time, modeled after Zerkle *et al.* (2005) with data updated for Mo (Anbar *et al.*, 2007; Scott *et al.*, 2008), Fe (Canfield *et al.*, 2008) and surface ocean O₂ (Holland, 2006).

As a consequence, prokaryotes should respond differently than eukaryotes to changes in metals and N budgets. It has therefore been suggested that the diversification of eukaryotes in marine ecosystems ~1.25 Ga was dictated by a shift from Mo-poor to Mo-rich oceans at that time (Anbar & Knoll, 2002).

In this paper, we summarize present knowledge of the metal requirements for N assimilation, and their biochemical basis, in both eukaryotes and prokaryotes. For each N assimilation pathway, we discuss the metals involved in the enzymes performing the reaction and discuss phylogenetic evidence for its evolutionary origin. We then synthesize this information with the emerging understanding of changes in marine metal availability over geologic time to develop hypotheses that may guide future research.


METALS AND N ASSIMILATION

Bioavailable nitrogen is found in four redox states in nature: +5 (NO₃⁻), +3 (NO₂⁻), 0 (N₂) and -3 (NH₄⁺, or in gaseous form, NH₃). To assimilate NO₃⁻, NO₂⁻ or N₂, electrons must be donated to reduce N to the -3 redox level of NH₄⁺, because only the most reduced form is used in biomolecules. In the following sections, we review the function and metal content of the enzymes responsible for the redox reactions involved in N assimilation: N₂ fixation (N₂ → NH₃), NO₃⁻ reduction (NO₃⁻ → NO₂⁻), NO₂⁻ reduction (NO₂⁻ → NH₄⁺) as well as of the enzymes used for assimilating DON and NH₄⁺ (summarized in Table 1). In addition, we summarize what is known about

the evolution of these enzymes based on gene sequences, which tells us whether eukaryotic proteins were inherited during endosymbiosis or whether they evolved separately in prokaryotes and eukaryotes. Analysis of such evolutionary relationships enables relative dating of N assimilation pathways and, hence, examination of the coevolution of metal availability in the oceans and photosynthetic N assimilatory enzymes.

Nitrogen fixation

Modern industry uses the Haber-Bosch process at extreme pressures and temperatures to fix N₂ into NH₃ over an Fe catalyst (Smil, 2004). Certain prokaryotes ('diazotrophs'), on the other hand, can perform this reaction at ambient pressures and temperatures. Heterotrophic and photosynthetic diazotrophs are prominent contributors to both the marine and terrestrial biospheres because they supply the major input of fixed N (Arrigo, 2005). The only oxygenic photosynthetic diazotrophs are cyanobacteria (Capone & Carpenter, 1982; Capone *et al.*, 1997; Stal & Zehr, 2008). Conversion of N₂ gas into NH₃ requires electrons, ATP and the metal-rich enzyme nitrogenase (abbreviated Nif), following the reaction:



The eight electrons required for each catalytic cycle are supplied by ferredoxin, a small [2Fe-2S]-cluster protein that accepts electrons from the photosynthetic electron transport chain (Schrautemeier & Bohme, 1985). The product NH₃ gas is rapidly converted into the soluble NH₄⁺ and directed into glutamine and glutamate biosynthesis (Wolk *et al.*, 1976; Meeks *et al.*, 1978; Carpenter *et al.*, 1992) (see Ammonium Assimilation section).

Nitrogenase is an oxygen-sensitive protein complex composed of two subunits: dinitrogenase ('Fe protein', or NifH) and dinitrogenase reductase ('Mo-Fe protein', or NifDK) (Berman-Frank *et al.*, 2003). Although cyanobacterial NifH has never been purified to homogeneity, cysteines that bind a [4Fe-4S] cluster from heterotrophic bacterial strains are conserved in cyanobacterial NifH sequences (Zehr *et al.*, 1997) (Fig. 2A). This suggests that cyanobacteria also contain a [4Fe-4S] cluster in NifH (Fig. 3A, Table 1). Electrons from reduced ferredoxin travel through this cluster to NifDK, which contains the Fe-Mo cofactor ('FeMoco'), the active site of N₂ reduction (Fig. 3A, Table 1). Numerous enzymes are involved in the biosynthesis of FeMoco, including two proteins, NifE and NifN, which serve as the scaffold upon which FeMoco is built (Rubio & Ludden, 2005, 2008).

The purified NifDK dimer from a freshwater cyanobacterium contains 2 Mo atoms and 20 Fe atoms (Hallenbeck *et al.*, 1979). This Fe content is lower than the ~25–38 Fe atoms found in NifDK dimers from heterotrophic bacteria (Howard & Rees, 1996; Kustka *et al.*, 2003a), although all Fe-S cluster-binding

Table 1 Characteristics of the key enzymes involved in N assimilation in phytoplankton

Enzyme	Metal co-factors	Substrates	Product	Distribution	Oxygenic photoautotrophs from which protein has been purified or overexpressed
Nitrogenase (Nif, 2-3 (?) subunits per enzyme)	4Fe-4S (NifH), 8Fe-8S, Mo(V/Fe)-7Fe-9S (NifDK)	N_2 , 8H^+ , 8e^- , 16ATP	2NH_3 , H_2 , 16ADP, 16P _i	Cyanobacteria	<i>Anabaena cylindrica</i> (Hallenbeck et al., 1979)
Ferredoxin-nitrate reductase (NarB)	4Fe-4S, molybdopterin guanine dinucleotide (Mo)	NO_3^- , 2e^- , 2H^+	NO_2^-	Cyanobacteria	<i>Plectonema boryanum</i> (Ida & Mikami, 1983; Mikami & Ida, 1984); <i>Synechococcus</i> sp. PCC 7942 (Rubio et al., 2002); <i>Aphanothece halophytica</i> (Thaivanich & Incharoensakdi, 2007)
NAD(P)H-nitrate reductase (NR, 4 subunits per enzyme)	Cytochrome <i>b</i> ₅₅₇ (Fe), sulfite-oxidase type molybdopterin (Mo)	NO_3^- , 2e^- , 2H^+	NO_2^-	Algae	<i>Chlorella vulgaris</i> (Solomonson, 1975); <i>Ankistrodesmus braunii</i> (de la Rosa et al., 1980, 1981)
Ferredoxin-nitrite reductase (NIR)	4Fe-4S, siroheme (Fe)	NO_2^- , 6e^- , 8H^+	NH_3	All	<i>Anabaena</i> sp. 7119 (Mendez & Vega, 1981); <i>Chlamydomonas reinhardtii</i> (Romero et al., 1987); <i>Phormidium laminosum</i> (Arizmendi & Serra, 1990)
Glutamine synthetase (GS)	Mn (?)	NH_4^+ , Glutamate, ATP	Glutamine ($\text{C}_2\text{H}_{10}\text{N}_2\text{O}_3$)	All	<i>Anabaena</i> sp. CA (Stacey et al., 1977); <i>Nostoc</i> sp. PCC 7120 (Orr et al., 1981); <i>Chlorella kessleri</i> (Sumar et al., 1984); <i>Anacystis nidulans</i> (Florencio & Ramos, 1985); <i>Chlorella sorokiniana</i> (Beudeker & Tabita, 1985); <i>Phormidium laminosum</i> (Blanco et al., 1989); <i>Calothrix</i> sp. PCC 7601 (Merida et al., 1990); <i>Synechocystis</i> sp. PCC 6803 (Merida et al., 1990; Garcia-Dominguez et al., 1997) <i>Prochlorococcus</i> spp. (El Alaoui et al., 2003)
Ferredoxin-glutamate synthase (GltS)	3Fe-4S	Glutamine ($\text{C}_2\text{H}_{10}\text{N}_2\text{O}_3$), 2-oxo-glutarate, 2e^- , 2H^+	2Glutamate ($\text{C}_5\text{H}_9\text{NO}_4$)	All (mainly used by cyanobacteria)	<i>Chlamydomonas reinhardtii</i> (Gotor et al., 1990); <i>Synechococcus</i> sp. PCC 6301 (Marques et al., 1992); <i>Synechocystis</i> PCC 6803 (Ravasio et al., 2002)
NADH-glutamate synthase (GltBD)	4Fe-4S (GltD), 4Fe-4S, 3Fe-4S (GltB)	Glutamine ($\text{C}_2\text{H}_{10}\text{N}_2\text{O}_3$), 2-oxoglutarate, 2e^- , 2H^+	2Glutamate ($\text{C}_5\text{H}_9\text{NO}_4$)	Some cyanobacteria, mainly used by eukaryotes	<i>Chlamydomonas reinhardtii</i> (Marquez et al., 1984)
Urease (Ure, usually 3 subunits per enzyme)	2Ni (di-nuclear)	Urea	NH_3 , H_2CO_3	Cyanobacteria/red algae	<i>Spirulina maxima</i> (Carvajal et al., 1982); <i>Anabaena doliolum</i> (Rai, 1989); <i>Anabaena cylindrica</i> (Argall et al., 1992); <i>Leptolyngbya boryana</i> (Jahns et al., 1995); <i>Anabaena</i> sp. PCC 7120 (Jahns et al., 1995); <i>Synechococcus</i> sp. WH 7805 (Collier et al., 1999)
Urea amidolyase	–	Urea	Allophanate	Green algae	No purification has been published.
Allophanate hydrolyase	–	Allophanate	NH_3	Green algae	<i>Chlamydomonas reinhardtii</i> (Maitz et al., 1982)
Cu-amino acid oxidase (CuAO, 2 subunits per enzyme)	Cu (mononuclear)	Amines (RCH_2NH_2), O_2	NH_3 , H_2O_2 , RCHO	Dinoflagellates/coccolithophores/cyanobacteria/green algae	No purification has been published.




Fig. 2 Schematic of N assimilatory protein alignments. Each horizontal shaded gray square represents a sequence of amino acid that folds into a protein. Each vertical colored bar represents the position of a conserved (present in the same position in the same proteins from different organisms) metal-binding amino acid in the protein sequence. It is clear that not all proteins are present in all organisms. For example, eukaryotes lack nitrogenase and possess different nitrate reductase proteins than prokaryotes, and most green algae do not possess urease nor copper-containing amine oxidase. The metal cofactors that are bound to the amino acids are shown above the sequences. Note the scale bar of 100 amino acids. Cysteines (abbreviated with a small 'C') are shown in yellow, histidines ('H') in red, serines ('S') in blue, lysines ('K') in dark green, aspartates ('D') in light green and glutamates ('E') in purple. See text for protein abbreviations. See Supporting Information (Supplementary text for Fig. 2) for details of the construction of the figure and for model organism species names.




Fig. 3 N assimilation pathways plotted versus redox state of N with metal cofactors colored (Fe: red, Mo: blue, Ni: green, Cu: purple, yellow atoms are S atoms), including N_2 fixation (a), prokaryotic NO_3^- reduction (b), eukaryotic NO_3^- reduction (c), NO_2^- reduction (d), urea hydrolysis (e) and amine oxidation (f). All assimilation pathways lead to NH_4^+ , which is assimilated following the pathways in Fig. 4. Enzyme and cofactor abbreviations are labeled and the N atom being assimilated is highlighted in teal in each pathway. AA stands for amine acid. Numbers in parentheses (i.e. x2) stands for the number of subunits in each enzyme.

cysteine residues align to identical positions in cyanobacteria (Fig. 2B,C). The ratio of NifH:NifDK in cyanobacteria is higher (3 : 1) than observed in other bacteria (2 : 1) (Reade *et al.*, 1999), suggesting that the overall Fe requirement may be similar between different organisms. The high Fe requirement for marine N_2 fixation is evident from experiments that have shown a steep drop-off of growth rate and nitrogenase activity below ~ 1 nM for *Trichodesmium* sp. (Berman-Frank *et al.*, 2001). Further biochemical characterization of nitrogenase proteins from marine diazotrophic cyanobacteria is crucial to characterizing the metal quotas of photosynthetic N_2 fixation in the sea.


The importance of Mo to N_2 fixation has been well-established since the 1930s (Bortels, 1930; Bortels, 1940). Nitrogenases that utilize alternatives to Mo were discovered more recently (Burns *et al.*, 1971; Bishop *et al.*, 1980). These alternative nitrogenases are distinct proteins encoded by separate genes (Robson *et al.*, 1986; Chisnell *et al.*, 1988; Pau *et al.*, 1989), which are expressed when Mo concentrations fall below 100 nM (Joerger & Bishop, 1988). Both vanadium-

Fe (Vnf) and Fe-Fe (Anf) nitrogenases exist (Eady, 1996); however, only three species of cyanobacteria contain *vnf* genes. All three known Vnf-containing species are strains of the freshwater cyanobacterium *Anabaena* (Kentemich *et al.*, 1988; Thiel, 1993; Boison *et al.*, 2006). No alternative nitrogenase genes have been found in the genomes of marine cyanobacteria, and a BLAST search of the J. Craig Venter Global Ocean Sampling database with the *Anabaena variabilis* ATCC 29413 VnfD protein (587 amino acids) returns marine peptides with only 34% identity at most. These are likely NifD protein hits, as Nif and Vnf proteins share $\sim 30\%$ sequence similarity (Raymond *et al.*, 2004). No cyanobacteria have been found to contain *anf* genes.

Nitrate reduction

Most, though not all, phytoplankton can assimilate NO_3^- . Prokaryotes and eukaryotes express evolutionarily-distinct proteins (Stolz & Basu, 2002) for this purpose, but regardless both kinds of nitrate reductases require Mo. NO_3^- assimilation

has been extensively reviewed for cyanobacteria, algae and plants (Guerrero *et al.*, 1981; Berges, 1997; Campbell, 1999; Moreno-Vivian *et al.*, 1999). In all cases, NO_3^- assimilation begins with the two-electron reduction to NO_2^- :



The protein responsible for the reduction of NO_3^- to NO_2^- in cyanobacteria is nitrate reductase, abbreviated NarB. This enzyme, like nitrogenase, depends on the donation of electrons from ferredoxin for its activity (Hattori & Myers, 1967; Manzano *et al.*, 1976; Mikami & Ida, 1984; Hirasawa *et al.*, 2004). Biochemical studies of NarB have shown that this protein binds one [4Fe-4S] cluster and one molybdopterin cofactor with cysteines (Rubio *et al.*, 1999; Rubio *et al.*, 2002) (Figs. 2D, 3B, Table 1). Two electrons travel from ferredoxin through the [4Fe-4S] cluster to the Mo atom in molybdopterin (Fig. 3B), which is reduced from the 6+ to the 4+ redox state by a two-electron reduction. NO_3^- then binds to the reduced Mo and is reduced to NO_2^- , returning Mo to the 6+ state (Jepson *et al.*, 2004).

Eukaryotic nitrate reductase arose through convergent evolution to serve the same role as NarB. This enzyme, abbreviated as NR, utilizes nicotinamide adenine dinucleotide (phosphate) ((NAD(P)H)), an organic cofactor that accepts electrons from ferredoxin, as its electron donor (Zumft *et al.*, 1969). NR in algae differs from NarB in cyanobacteria, in that NR contains four identical protein subunits, each containing one atom of Mo, whereas NarB contains only one subunit. Each NR subunit contains three prosthetic groups: flavin adenine dinucleotide (FAD) (Zumft *et al.*, 1970), cytochrome b_{557} (Ahmed & Spiller, 1976), and molybdopterin of a slightly different structure than in NarB (Solomonson *et al.*, 1975; Solomonson *et al.*, 1986) (Fig. 3C, Table 1). Cysteines that bind molybdopterin and cytochrome in NR from yeast (Fischer *et al.*, 2005) and corn (Lu *et al.*, 1995) align to conserved position in NR from diatoms, and green and red algae with primary plastid symbionts derived from cyanobacteria (Fig. 2E), suggesting that the metal-containing cofactors of NR are likely conserved among these evolutionary distant organisms. During catalysis, electrons donated by NAD(P)H reduce FAD, which transfers the electrons to cytochrome b_{557} . Electrons then tunnel to the active site Mo atom in the molybdopterin, where NO_3^- is reduced to NO_2^- (Skipper *et al.*, 2001) (Fig. 3C, Table 1).

The Fe content of NR is well documented (Aparicio *et al.*, 1971; Cárdenas *et al.*, 1972). Four Fe atoms, corresponding to four cytochrome molecules, are present in all described algal NRs (de la Rosa *et al.*, 1981; de la Rosa, 1983; Solomonson *et al.*, 1986). Although the physiological requirement of Mo for nitrate reduction has been established for fungi (Steinberg, 1937; Nicholas & Nason, 1954; Nicholas *et al.*, 1954) and green algae (Arnon *et al.*, 1955; Aparicio *et al.*, 1971; Cárdenas *et al.*, 1971; Vega *et al.*, 1971), the exact number of atoms in

NR is still unclear. It was expected that each of the four NR subunits contain one molybdopterin cofactor, and thus one Mo atom, as has been found for NR purified from one green algal species (Solomonson *et al.*, 1986). However, another green algal species was found to contain only two Mo atoms in NR (de la Rosa *et al.*, 1981; de la Rosa, 1983). Further study is necessary to determine whether Mo content is in fact variable in eukaryotic NR.

NO_3^- reduction also occurs in anaerobic bacteria, which utilize NO_3^- as a terminal electron acceptor during dissimilatory NO_3^- reduction in anoxic waters and sediments. Dissimilatory nitrate reductases that contain no Mo or Mo-alternatives (heme Fe and V) have been discovered, but only in organisms from extreme environments (Antipova *et al.*, 1998; Antipov *et al.*, 2003, 2005).

Nitrite reduction


Following NO_3^- reduction, a six-electron reduction of NO_2^- to NH_4^+ is essential for N incorporation. NO_2^- assimilation can also occur independently from NO_3^- assimilation. Nitrite reductase (NiR) is an iron-rich protein, containing one [4Fe-4S] cluster and one heme (Fig. 3D, Table 1). The [4Fe-4S] cluster transfers electrons from ferredoxin to Fe in siroheme, where six electrons reduce NO_2^- to NH_4^+ (Kuznetsova *et al.*, 2004), following the overall reaction:



Biochemical and phylogenetic evidence suggests that NiR in algae was inherited from cyanobacteria. This inference follows from the finding that cysteine residues bound to the [4Fe-4S] cluster and heme in the crystal structure of spinach nitrite reductase (Swamy *et al.*, 2005) are conserved in the NiR sequences from cyanobacteria and algae (Fig. 2F). Hence, NiR's Fe cofactors appear to be bound in a similar fashion in all photoautotrophs. This suggests that the origin of NiR in algae may originally derive from the endosymbiont that lead to the modern plastid. In addition to their biochemical similarities, phylogenetic analysis of the NiR-encoding *nirA* gene shows that cyanobacteria group near the base of the tree, with higher plant and algal sequences branching out from the cyanobacterial clade (Wang *et al.*, 2000).

DON assimilation


Dissolved organic nitrogen (DON) occurs in many chemical forms; here, we discuss only urea and amino acid assimilation. The enzyme urease, found in cyanobacteria and most algae and plants, catalyzes the hydrolysis of urea, producing two molecules of ammonia and one molecule of carbonic acid (Fig. 2E, Table 1):



Urease isolated from cyanobacteria and diatoms contains three subunits, usually two small and one large (Fig. 3E) (Carvajal *et al.*, 1982; Rai, 1989; Argall *et al.*, 1992; Jahns *et al.*, 1995; Collier *et al.*, 1999; Palinska *et al.*, 2000). Nevertheless, the same amino acids (four histidine residues, one lysine and one arginine) that bind Ni in the large subunit of urease from the heterotrophic bacterium *Klebsiella aerogenes* (Jabri *et al.*, 1995) are conserved in cyanobacterial and diatom ureases (Fig. 2G) suggesting that Ni plays a similar role in the ureases of these species as in other bacteria. Addition of Ni is essential for the growth of coastal diatoms (Oliveira & Antia, 1984, 1986a,b; Egleston & Morel, 2008) and marine cyanobacteria *Synechococcus* (Dupont *et al.*, 2007) on urea as the sole N source. Urease activity has been shown to increase with rising Ni concentration (Mackerras & Smith, 1986; Price & Morel, 1991). Urea assimilation represents one of the few biological uses of Ni in photoautotrophs, besides Ni-superoxide dismutase (Dupont *et al.*, 2007) and hydrogenase (Tamagnini *et al.*, 2002). Interestingly, an alternative urea assimilation pathway exists in certain green algae, which does not require Ni for urea assimilation (Rees & Bekheet, 1982). Instead, ATP is used in a two-step hydrolysis of urea.

The relative age of urease is unknown. This enzyme is member of an ancient and diverse protein superfamily with conserved 3-D structure (Holm & Sander, 1997), but the timing of urease evolution within this family remains unclear. The phylogeny of *ureC*, which encodes the large, Ni-binding subunit of urease, suggests that cyanobacteria and eukaryotic *ureC* genes evolved independently after an early gene duplication event (Koper *et al.*, 2004). Very little phylogenetic work has investigated alternative non-Ni containing ureases in green algae (Syrett & Alhouly, 1984). These proteins deserve further research attention, as they may afford an evolutionary advantage for more efficient green algal assimilation of urea in low-Ni aquatic systems.

Algae can also assimilate N from free amino acids by the action of a Cu-containing cell-surface amine oxidase (CuAO) (Fig. 3F). This is the only N assimilatory protein yet discovered that works at the outside surface of the cell, and functionally parallels alkaline phosphatase, the enzyme that strips phosphate groups from nucleic acids during P assimilation. CuAO oxidizes amino acids to produce hydrogen peroxide (H₂O₂), α -keto acids and extracellular ammonia, following the reaction:



The NH₃ is subsequently taken up through specific transporters and assimilated by the organism (see Ammonium Assimilation section). Amine oxidase activity has been observed for phytoplankton with secondary plastid symbionts derived from red algae including coccolithophorids and dinoflagellates (Palenik & Morel, 1990a,b, 1991). Although the Cu content of algal CuAO has not been confirmed, it is likely that they

resemble other amine oxidases, which contain two atoms of Cu (one in each of the two subunits) because experiments have shown that growth of the marine coccolithophorid *Pleurochrysis carterae* on amines requires 10 nM Cu (Palenik & Morel, 1991). However, no other studies looking at the effect of amine sources on CuAO activity or intracellular Cu of algae have yet been performed. The crystal structure of pea CuAO shows that Cu is coordinated at the active site by three histidine residues (Kumar *et al.*, 1996). Histidine residues are conserved in phytoplankton CuAO as predicted by the gene sequence of green algae (Fig. 2H), suggesting that the active site, and perhaps the metal requirement, of algal CuAO may be similar to that of higher plants. CuAO is present in numerous eukaryotes (Cona *et al.*, 2006), but only a small fraction of prokaryotic genome sequences analyzed contain AO-encoding genes (Andreini *et al.*, 2008). However, the genomes of numerous marine and freshwater cyanobacteria contain putative CuAO-encoding genes. Further phylogenetic and biochemical characterization of this protein is essential to evaluate its Cu content and thus the ecological and evolutionary consequences of this pathway.

Ammonium assimilation

The last step in N assimilation is conversion of NH₄⁺ to glutamate. Regardless of the N source, all N assimilation pathways end with NH₄⁺ assimilation through the glutamine synthetase-glutamate synthase (GS-GltS) pathway, also known as the GS-GOGAT pathway (Muro-Pastor *et al.*, 2005). First, NH₄⁺ is incorporated into glutamate, yielding glutamine, and then N is transferred as an amide group to 2-oxoglutarate, producing an additional molecule of glutamate to be used for amino acid synthesis (Fig. 4A,B, Table 1). The first enzyme in the GS-GltS pathway is the ATP-dependent enzyme glutamine synthetase (GS), which catalyzes the formation of glutamine (Fig. 4A, Table 1):



GS is most often a large, multi-subunit protein (Stacey *et al.*, 1977; Sampaio *et al.*, 1979; Orr *et al.*, 1981; Sumar *et al.*, 1984; Beudeker & Tabita, 1985; Florencio & Ramos, 1985; Merida *et al.*, 1990; Yuan *et al.*, 2001; El Alaoui *et al.*, 2003) present in thylakoid membranes (Lopez-Ruiz *et al.*, 1991) and requires divalent cations, such as Mg or Mn, for activity (Blanco *et al.*, 1989). The wide distribution of GS-encoding genes throughout the three domains of life suggests that this gene is ancient and has likely been horizontally transferred between the bacterial and archeal domains (Raymond, 2005).

The second enzyme in the GS-GltS pathway is glutamate synthase (GltS, also known as GlsF or GOGAT) (Suzuki & Knaff, 2005). GltS catalyzes the conversion of glutamine and 2-oxoglutarate (a product of the tricarboxylic acid cycle (TCA cycle)) into two moles of glutamate, via a 2-electron reduction:





Fig. 4 The NH_4^+ assimilation pathway in primary producers with Fe colored in red and S colored in yellow. NH_4^+ taken up from the environment or reduced via the pathways in Fig. 3 is converted to glutamine (A) and subsequently to two molecules of glutamate with either ferredoxin (B) or NADH (C) supplying electrons for this process. One molecule of glutamate is recycled to make sure glutamine and the other is converted into biomolecules requiring N, such as other amino acids, heme or chlorophyll.



In cyanobacteria, electrons for this process travel from ferredoxin through a [3Fe-4S] cluster in GltS to the organic molecule flavin mononucleotide (FMN) at the active site of glutamate formation (Merida *et al.*, 1990; Marques *et al.*, 1992; Navarro *et al.*, 2000; Ravasio *et al.*, 2002) (Fig. 4B). The crystal structure of GltS from *Synechocystis* PCC 6803 reveals that the [3Fe-4S] cluster is coordinated by three cysteines (van den Heuvel *et al.*, 2002, 2003), which are conserved in GltS sequences from other cyanobacterial and algae (Fig. 2I), supporting the essential role for Fe in phytoplankton glutamate synthesis. However, although algae possesses genes encoding GltS-like cyanobacteria, many express a larger two-protein complex, abbreviated GltBD, for glutamine assimilation (Cullimore & Sims, 1981; Marquez *et al.*, 1984) (Fig. 4C, Table 1). Like NR (see Nitrate Reduction section), GltBD accepts electrons from NADH instead of ferredoxin (Clayton & Ahmed, 1986). Sequence similarity to the crystallized subunit GltB of the greater GltBD complex from a diazotrophic bacterium (*Azospirillum brasilense*) (Binda *et al.*, 2000) (Fig. 2I),

suggests that algal GltB also binds one [3Fe-4S] cluster. The additional subunit (GltD) in *A. brasilense* contains two [4Fe-4S] clusters (Vanoni & Curti, 1999; Vanoni *et al.*, 2005) (Table 1), and the cysteines that bind these clusters are conserved in algal GltB sequences (Fig. 2J). Thus, GltBD contains 11 Fe atoms per enzyme complex, compared to only 3 in GltS. Why would utilizing NADH instead of ferredoxin as an electron donor for glutamate synthesis require a nearly fourfold increase in Fe? This evolutionary question remains unanswered.

Phylogenetic analyses suggest that GltS-encoding genes are ancient, perhaps already present in the last universal common ancestor, and have diversified to use different electron donors (Raymond, 2005). The *gltS* gene (also known as *glsF*) has been found encoded in the chloroplast genome of numerous primary red algae, suggesting that this gene, like *nirA*, had an endosymbiotic origin (Valentin *et al.*, 1993). However, this gene is not present in the chloroplast genome of primary green algae, suggesting either an early transfer to the nucleus or a nonendosymbiotic origin (Grzebyk *et al.*, 2003). Despite these differences, both primary red and green algae now rely primarily on Fe-rich GltBD enzymes for NH_4^+ assimilation (Clayton & Ahmed, 1986).

N ASSIMILATION AND METALS THROUGH EARTH HISTORY: CO-EVOLUTION AND COMPETITION

The redox state of the oceans has changed dramatically through time. The O₂ content of seawater has, in turn, likely influenced the speciation and availability of surface and deep-water metal concentrations (Fig. 1). Our understanding of these ancient variations is developing rapidly, particularly as regards the abundances of Fe and Mo (Rouxel *et al.*, 2005; Anbar & Rouxel, 2007; Anbar *et al.*, 2007; Scott *et al.*, 2008), two metals that are important in N assimilation enzymes. The relationships between O₂, N speciation and metal availability create selection pressures that could have shaped the evolution of metals in N assimilation enzymes. If so, the metal requirements of N assimilation pathways found in organisms today may be a legacy of evolution in ancient oceans (Saito *et al.*, 2003; Morel, 2008).

We consider this possibility in the succeeding discussion, focusing in particular on Fe and Mo. Aside from their importance in N assimilation, these metals are especially interesting in an evolutionary context because their ocean abundances change in an opposite sense to each other in response to changes in ocean oxygenation (Fig. 1). Continental Mo sulfides are only weathered into rivers in the presence of O₂, whereas Fe hydrolyzes into insoluble Fe oxides. Hence, we propose a sequence for the co-evolution of Fe- and Mo-containing N assimilation pathways cognizant of historical changes in ocean metal availability and environmental redox conditions during approximately the first half of Earth history. We then explore the implications for the ecological competition between prokaryotes and eukaryotes in the face of subsequent changes in ocean oxygenation.

Co-evolution of N assimilation and metal availability

The early oceans contained a very different suite of bioavailable elements than today. The most primitive N assimilatory enzymes would have evolved to assimilate reduced N species, instead of the oxidized N present in the modern environment (Falkowski & Godfrey, 2008). In order to perform redox chemistry, metals were required. Fe, in the form of Fe-S clusters and heme, was both bioavailable and catalytically useful for these reactions. As O₂ began to rise, new enzymes utilizing more oxidized N substrates and metals made more bioavailable by oxygenation evolved.

The anoxic Archean


Molecular O₂ was absent or scarce in the Archean atmosphere and oceans (Holland, 2006). Under these conditions, the most abundant form of bioavailable N was probably NH₄⁺, rather than NO₃⁻ as today (Boyd & Philippot, 1998; Beaumont & Robert, 1999; Papineau *et al.*, 2005). Some N may also have been present as NO₂⁻. However, then, as now, the vast majority of N in the oceans and atmosphere was present as N₂ (Kasting & Catling, 2003), which is not readily bioavailable.

Before the evolution of biological N₂ fixation, the supply of fixed N to the oceans would have been limited by the rate of abiotic N₂ fixation. This rate is unknown, but could have been significant. As in biology, metal catalysts were probably important for abiotic N₂ fixation. For example, iron-sulfide minerals likely fixed some N₂ into NH₄⁺ at Archean hydrothermal vents (Schoonen & Xu, 2001). Some N₂ was probably also fixed without metals, by atmospheric reactions driven by lightning. These reactions convert N₂ into nitric oxide (NO), which dissolves into the ocean as precipitable HNO₂ and HNO₃ (Yung & McElroy, 1979; Mancinelli & McKay, 1988; Navarro-González *et al.*, 1998; Navarro-González *et al.*, 2001). Bolide impacts and subaerial volcanoes could have supplied additional fixed N as N-oxides (Mancinelli & McKay, 1988; Mather *et al.*, 2004). In the ocean, NO₂⁻ could have been abiotically reduced to NH₄⁺ by abundant ferrous Fe (Summers & Chang, 1993).

Thus, it is reasonable to hypothesize that both NH₄⁺ and NO₂⁻ were present in the Archean oceans and hence that the first N assimilatory enzymes to evolve were those involved in NO₂⁻ and NH₄⁺ assimilation: nitrite reductase (NiR) and glutamine synthetase (GS)/glutamate synthase (GltS(BD)), respectively (Fig. 5). The genetic sequence, protein structure and cofactor assemblage of these enzymes are highly conserved across all photoautotrophs (Luque *et al.*, 1993; Valentin *et al.*, 1993; Raymond, 2005), supporting their great antiquity. Further evidence of their early evolution comes from phylogenetic evidence suggesting that NiR and GltS proteins were transferred from cyanobacteria to a eukaryotic host during endosymbiosis (Temple *et al.*, 1998; Wang *et al.*, 2000). For these reasons, the early evolution of GS and GltS(BD) has previously been proposed by Papineau *et al.* (2005).

The early Fe requirement for assimilation of NO₂⁻ and NH₄⁺ through the action of Fe-containing NiR and GltS(BD) proteins is fully consistent with our understanding of metal availability in Archean oceans. The high Fe content of Archean banded iron formations suggests that Fe was present in micromolar concentrations in the Archean oceans, compared to sub-nanomolar quantities in the modern sea (Holland, 1973). Iron isotopic compositions from sulfides in black shales paint a picture of high Fe content in the Archean oceans compared to the oceans of the Proterozoic and Phanerozoic (Kump, 2005; Rouxel *et al.*, 2005; Johnson *et al.*, 2008). Fe content may have fluctuated in response to relative input from hydrothermal vents versus export as Fe oxides in banded iron formations (Rouxel *et al.*, 2005). The question of whether N assimilation first evolved at seafloor hydrothermal systems or in the open ocean remains unanswered. The Archean oceans were low in S (Canfield *et al.*, 2000), however, NiR and GltS(BD) contain Fe-S clusters, so evolution at mineral-water interfaces has some compelling logic. It is also clear that oxyanions such as Mo and V are absent from NiR and GltS(BD), consistent with accumulating evidence of very low concentrations of these metals in the Archean ocean (Anbar *et al.*, 2007; Scott *et al.*, 2008).

Fig. 5 Trends in the relative abundance of metals involved in N assimilation (Fe, Mo, Ni and Cu) and N species (N_2 , NO_3^- , NO_2^- , NH_4^+ urea and amino acids) through time. Color darkness represents relative average seawater concentration, with darker colors representing higher concentrations, and lighter colors representing lower concentrations. Note that very little data exist on the abundance of organic N (urea and amino acids) through time. Solid blue bars indicate periods when it has been fairly well-established that N assimilation enzymes had evolved, whereas question marks indicate periods when they may have existed. Anf = Fe-Fe nitrogenase; Vnf = V-Fe nitrogenase; Nif = Mo-Mo nitrogenase, NarB = cyanobacterial nitrate reductase; NR = eukaryotic nitrate reductase; NiR = nitrite reductase; Ure = urease; AO = amine oxidase.



Abiotic production of NH_4^+ and NO_2^- was probably not sufficiently rapid to support a biosphere comparable to today's, creating a selection pressure that favored the evolution of biological N_2 fixation pathways (Navarro-Gonzalez *et al.*, 2001). Fossil and phylogenetic evidence suggest that some cyanobacteria had acquired the ability to fix N_2 by the early Paleoproterozoic (Tomitani *et al.*, 2006), but there is reason to suspect that N_2 fixation evolved much earlier. The high similarity between *nifD*, *nifK*, and *nifE* and *nifN* genes scattered throughout bacteria and archaeal genomes implies that these genes duplicated before the last universal common ancestor, presumably early in the Archean (Fani *et al.*, 2000). This suggested antiquity is consistent with the O_2 -sensitivity of all forms of nitrogenase, which implies that they evolved before O_2 was pervasive in the environment.

The existence of three forms of nitrogenase using different metals (Mo, V and Fe) has led to the suggestion that Anf – the Fe-Fe nitrogenase – evolved in the Archean, when Fe was plentiful but before Mo (and V) was present in any significant quantities (da Silva & Williams, 1991) (Fig. 1). Intriguingly, if Anf is ancestral it is conceivable that this enzyme complex developed for reasons other than N_2 fixation. Anf is the least effective nitrogenase of the three, but is the most effective at H_2 production (Schneider *et al.*, 1997). Thus, Anf may have initially evolved in Fe-rich oceans for H_2 production and subsequently been adopted for nitrogenase function (Silver & Postgate, 1973; Normand & Bousquet, 1989; Fani *et al.*, 2000). However, the phylogenetic record does not present unambiguous evidence of an evolutionary sequence among the three

alternative nitrogenases, probably because of lateral gene transfer and gene loss (Raymond *et al.*, 2004). Thus, although the high Fe content of Anfs is consistent with Archean geochemistry, its place in evolutionary history is obscure.

The oxygen transition

Oxygenic photosynthesis may have evolved by 2.7 Ga (Brocks *et al.*, 1999, 2003a,b; Eigenbrode & Freeman, 2006), based on the interpretation of molecular biosignatures and stromatolite paleoenvironments (Buick, 1992, 2008), but these interpretations are not undisputed (Kirschvink *et al.*, 2000; Kirschvink & Kopp, 2008; Rasmussen *et al.*, 2008). However, few dispute that by 2.3 Ga, O_2 in the atmosphere rose above $\sim 10^{-5}$ PAL (Holland, 2006). The atmosphere and at least the surface oceans were pervasively oxidized by 1.8 Ga, whereas the deep sea remained euxinic (Fig. 1) (Poulton *et al.*, 2004). This rise of O_2 would have stabilized NO_2^- and then NO_3^- in the ocean (Fig. 5) (Beaumont & Robert, 1999; Papineau *et al.*, 2005).

The reason for the delay of at least 300 Myr between the origin of oxygenic photosynthesis and the oxygenation of the environment, assuming it occurred, is unclear. Several processes could have conspired to maintain a low- O_2 condition (Holland, 2002; Catling & Claire, 2005; Fennel *et al.*, 2005; Kump & Barley, 2007). Among these were weathering reactions with iron sulfide (pyrite) and other reduced minerals, some of which are host phases of Mo and other bioessential metals. Once mobilized by oxidative weathering, Mo was stabilized in oxygenated waters as the oxyanion molybdate (MoO_4^{2-}), making it widely available for the first time for use

in N-cycle enzymes such as Nif and nitrate reductase (cyanobacterial NarB and eukaryotic NR). Hence, the use of Mo in N-cycle enzymes probably evolved during a time when biogenic O₂ was commonly produced but not yet pervasive.

During this period, the residence times of both O₂ and Mo in the oceans were short, and hence their concentrations variable. Therefore, it is possible to imagine that both Fe and Mo were sporadically available in near-shore environments before the rise of a more thoroughly oxygenated environment. Recent geochemical findings suggest that such conditions did, indeed, occur after 2.7 Ga in the form of at least one ‘whiff’ of O₂ to p.p.m. levels, and associated marine Mo enrichment, at 2.5 Ga (Anbar *et al.*, 2007). It is even possible that such events were common between 2.7 and 2.3 Ga but never predominant as Mo concentrations in late Archean shales are generally very low (Cameron & Garrels, 1980; Davy, 1983; Yamaguchi, 2002).

It was presumably during this time that Nif became the dominant N₂ fixation enzyme in aerobic microbial communities as a result of the rise of marine Mo concentrations (Figs 1, 5) (Anbar *et al.*, 2007; Scott *et al.*, 2008). This hypothesis is consistent with the N isotopic data of Garvin *et al.* (2009). The key advantage of Mo is that Nif is ~15 times more efficient at *in vivo* N₂ fixation than the alternative forms of nitrogenase (Schneider *et al.*, 1997). Sporadic availability of both Mo and Fe would have facilitated the incorporation of Mo at the enzyme active site while still maintaining heavy dependence on Fe for the rest of the Fe-S clusters. Numerous microbes, namely methanogenic Archaea and anoxygenic photosynthetic Bacteria (likely the predecessors of cyanobacteria (Blankenship, 1992, 2001; Burke *et al.*, 1993)), possess all three sets of nitrogenase genes (Madigan, 1995; Masepohl *et al.*, 2004; Oda *et al.*, 2005), allowing them to inhabit both high- and low-Mo environments. Intriguingly, Anf in some anoxygenic photoautotrophs can incorporate a Mo-Fe cofactor when Mo is available (Gollan *et al.*, 1993). The ability to repurpose a different cofactor for the same gene product suggests that primitive nitrogenases were more flexible in metal use, consistent with evolution in a world with variable Fe and Mo availability.

Nitrate reductase enzymes may have also evolved during this period of low, possibly transient, levels of O₂ and Mo in the environment (Garvin *et al.*, 2009). The assimilatory form of prokaryotic nitrate reductase shares a common ancestor with the dissimilatory (or respiratory) nitrate reductase utilized in the first step of denitrification (Stolz & Basu, 2002). If denitrification evolved before NO₃⁻ assimilation, it is possible to constrain the timing based on geologic evidence. Isotopic signatures of denitrification are found as early as 2.5 Ga (Garvin *et al.*, 2009), providing a minimum age for the origin of this metabolism. These signatures are also seen in ~2.1 Ga sediments (Boyd & Philippot, 1998; Beaumont & Robert, 1999; Papineau *et al.*, 2005).

However, the prokaryotic nitrate reductase family most likely evolved much earlier than 2.1 Ga. Phylogenetic evidence

suggests an origin of the gene family before the divergence of eubacteria and archaea (Petri & Imhoff, 2000; Philippot, 2002) between 3 and 4 Ga (Feng *et al.*, 1997), with subsequent passage of the gene among prokaryotes through horizontal gene transfer (Stolz & Basu, 2002). The apparent antiquity of nitrate reductase is puzzling, given that sources of NO₃⁻ and Mo would have been limited in the anoxic Archean. It is possible that different metals – perhaps V and Fe – were utilized in place of Mo in early nitrate reductases, but few modern examples of alternative nitrate reductases exist (see Nitrate Reduction section). Alternatively, the antiquity of NarB may tell us that slightly oxidizing conditions, such as those documented at 2.5 Ga (Anbar *et al.*, 2007; Kaufman *et al.*, 2007), occurred episodically through the Archean.

Competition between prokaryotes and eukaryotes in metal-starved marine ecosystems

Once the N assimilation metabolisms became established, the evolutionary focus likely shifted increasingly toward competition between marine organisms for dominance. Cyanobacteria prevailed during the first three quarters of Earth history, whereas algae rose to prevalence in the Phanerozoic (Falkowski *et al.*, 2004; Knoll, 2007). In this section, we discuss the progression of marine primary producers with regard to metal requirements for N assimilation.

After the first oxygenation event

The Great Oxidation Event (GOE) ushered in a new period of the Earth’s chemical history after 2.3 Ga. Paradoxically, this dramatic rise in atmospheric O₂ likely led to intensified anoxia and the generation of H₂S in deep ocean waters (Canfield, 1998). This counterintuitive consequence arises from the enhanced influx of sulfate as a result of oxic weathering of sulfide minerals on the continents, stimulating the action of sulfate-reducing bacteria in the oceans (Canfield, 1998). If the partial pressure of O₂ in the atmosphere was still significantly lower than today’s – as generally believed – then the rate of H₂S generation could have exceeded the rate of O₂ supply to deep waters in many places, resulting in an ‘H₂S maximum’ analogous to the ‘O₂ minimum’ in modern oceans. This hypothesis is supported by mounting evidence for widespread deep-water anoxia and high sulfide (‘euxinia’) by 1.84 Ga (as compared to today), persisting until the Neoproterozoic (Shen *et al.*, 2003; Arnold *et al.*, 2004; Poulton *et al.*, 2004; Brocks *et al.*, 2005; Scott *et al.*, 2008).

The advent of this euxinic ‘Canfield Ocean’ would have had important consequences for metal availability and hence for N assimilation. Sulfide in the water column would have facilitated rapid removal of Mo (and some other sulfide-reactive transition metals) to sediments, limiting the accumulation of Mo in the oceans despite the enhanced influx from oxic continental weathering. At the same time, the rise of surface-water oxygen coupled with deep-water sulfidic conditions would have

Table 2 Comparison of the specific activities of prokaryotic (P) to different eukaryotic (E) nitrate reductase enzymes

Specific activity ($\mu\text{moles NO}_2^-$ formed $\text{min}^{-1} \text{mg enzyme}^{-1}$)	Species	Reference
20–40	<i>Synechococcus</i> PCC 7942 (cyanobacteria, P)	Rubio <i>et al.</i> (2002)
72–80	<i>Ankistrodesmus braunii</i> (green alga, E)	de la Rosa <i>et al.</i> (1981)
83–86	<i>Chorella vulgaris</i> (green alga, E)	Solomonson <i>et al.</i> (1975)
2050	<i>Thalassiosira pseudonana</i> (diatom, E)	Amy & Garrett (1974)

precipitated a crash in Fe concentrations compared to those common in the Archean (Fig. 1).

Under these conditions, N_2 -fixing prokaryotes would have struggled to obtain sufficient Mo and Fe for Nif and Anf, potentially resulting in persistent N-limitation in the oceans (Anbar & Knoll, 2002). Anf and Vnf do not exist in marine cyanobacteria today, so it is possible that Mo was absolutely necessary for cyanobacterial N_2 fixation throughout Earth's history. Experiments testing this hypothesis have shown that the rates of growth and N_2 fixation by certain cyanobacteria drop when the concentration of Mo is $<5 \text{ nM}$ and when Fe is less than $<1 \text{ nM}$ in the growth media (Berman-Frank *et al.*, 2001; Zerkle *et al.*, 2006). Measurements of Mo in ancient sediments together with simple mass balance models suggest that Mo concentrations were likely 10–20 nM in the Proterozoic, and even lower in the Archean (Anbar *et al.*, 2007; Scott *et al.*, 2008). Combined, these experiments and observations suggest that Mo availability in mid-Proterozoic oceans was close to the levels that are problematic for N_2 fixation by extant cyanobacteria, and at or below that level in the Archean.

Would Proterozoic oceans have contained sufficient Mo for NO_3^- assimilation? We sought to determine the cyanobacterial Mo requirements for NO_3^- assimilation by growing the filamentous, heterocystous cyanobacterium *Nostoc* sp. PCC 7120 with NO_3^- as the sole N source, with (1500 nM) and without ($\sim 0.5 \text{ nM}$) Mo in semi-continuous batch culturing experiments to maintain a growth rate of ~ 1 doubling per day. After 1 month, we found no difference in growth rate or intracellular N between Mo-deplete and replete cultures (Fig. 6). It appears, then, that cyanobacteria may be specially adapted to grow on NO_3^- with minimal Mo. This could be accomplished through the effective uptake of low quantities of Mo by high-affinity molybdate-uptake systems (Thiel *et al.*, 2002; Zahalak *et al.*, 2004), long-term Mo storage, or expression of nitrate reductases containing alternative metal centers (see Nitrate Reduction section). Thus, cyanobacteria could have used NO_3^- as an alternative N source if N_2 fixation was Mo-limited in the mid-Proterozoic. Cyanobacteria were the major marine primary producers throughout the Proterozoic (Knoll, 2007), perhaps in part resulting from their ability to assimilate NO_3^- even at low Mo concentrations.

Phylogenetic evidence suggests that NR arose soon after the origin of eukaryotes (Stolz & Basu, 2002) in the late Archean (Brocks *et al.*, 1999; Hedges *et al.*, 2001; Brocks *et al.*, 2003a,b). The Mo requirement for eukaryotic nitrate reductase (NR)




Fig. 6 Results of an experiment in which the heterocystous cyanobacterium *Nostoc* sp. PCC 7120 was grown for 1 month and diluted every 3–5 days with new media containing NO_3^- (15 mM) and either no Mo ('-Mo', blank was 0.5 nM) or 1500 nM Mo as sodium molybdate. (A) and (B) Photographs of one replicate culture bottle from each condition with Mo (A) and without Mo (B) on the final day of the experiment. (C) Physiological measurements taken on the final day of the experiment. Intracellular Mo was $\sim 4\times$ higher in the +Mo bottle than the -Mo bottle, but this intracellular difference had no effect on the growth rate (dark circles), nor the intracellular N content of the cultures (open circles). Chlorophyll concentrations were 1.0 mg L^{-1} in both conditions on the last day of the experiment. These results show that cyanobacteria are adapted to cope with very low environmental Mo levels when they are grown with a source of nitrate. See Supporting Information (Supplementary text for Fig. 6) for details about the growth conditions of this experiment.

may be considerably higher than that of the cyanobacterial NarB enzyme. NR requires four times as much Mo per enzyme as NarB per enzyme complex, as a result of NR's tetrameric structure. However, NR is up to $100\times$ more active than the cyanobacterial NarB (Table 2), and thus the Mo requirement of NR may actually be lower than that of NarB. This alternative hypothesis is not supported by preliminary studies, which show that under low Mo, benthic bacteria actively assimilate and denitrify NO_3^- whereas phytoplankton (dominated by algae) show little NO_3^- uptake (Axler & Reuter, 1996). Assuming that the Mo requirement for NR is higher than that of NarB, eukaryotic growth may have been N-limited in the Mesoproterozoic ocean before the major increase in marine Mo $\sim 0.6 \text{ Ga}$ (Scott

et al., 2008; Wille *et al.*, 2008) especially because eukaryotes lack the ability to directly utilize N_2 . The scenario of Mo limitation of eukaryote proliferation fits well with the fossil record of algae which displays a radiation of green algae in the Cambrian (Revill *et al.*, 1994; Knoll, 2007). These green algae could have fueled the food web that emerged after the Cambrian Explosion.

The transition to the (mostly) oxic Phanerozoic

Neoproterozoic marine life was influenced by widespread glaciations, the recurrence of banded iron formations and finally the oxygenation of the deep sea between ~0.58–0.56 Ga (Fike *et al.*, 2006; McFadden *et al.*, 2008; Scott *et al.*, 2008; Shen *et al.*, 2008). Ferruginous anoxic deep-water conditions may have prevailed from ~0.7–0.55 Ga (Fig. 1) (Canfield *et al.*, 2008) and could have alleviated Fe stress during upwelling events. At the same time (~0.6 Ga), Mo concentrations were rising (Fig. 1) (Scott *et al.*, 2008). Concurrent Fe and Mo fertilization of surface waters could have fueled primary productivity in the oceans, supplying food with a high N content to nourish early multicellular organisms in the Ediacaran ocean.

Scattered throughout the early Phanerozoic were warm intervals when the ocean reverted back to deep-water anoxia or euxinia. These periods are thought to have been characterized by N limitation of primary productivity (Saltzman, 2005). Ocean anoxic events (OAEs) of the Paleozoic could have reduced Mo concentrations to 30–50% of modern levels (Algeo, 2004), and ocean Mo during Mesozoic OAEs may have been drawn down to as little as ~1% of modern levels (Anbar & Gordon, 2008; Pearce *et al.*, 2008). At the Permian-Triassic (P/T) boundary there are indications that euxinia reached the photic zone in much of the oceans (Grice *et al.*, 2005; Kump *et al.*, 2005), which may have vastly depleted transition metal concentrations. Algal growth and evolution may have been slowed during these periods (Falkowski *et al.*, 2004). Biomarker and isotopic evidence shows that cyanobacterial populations surged at the P/T boundary (Xie *et al.*, 2005) and during Cretaceous OAEs (Kuypers *et al.*, 2004; Ohkouchi *et al.*, 2006; Junium & Arthur, 2007; Kashiyama *et al.*, 2008), consistent with the hypothesis that cyanobacteria are particularly well-adapted to cope with metal draw-down in euxinic oceans. If these low-Mo periods limited eukaryotic N metabolism, cyanobacteria would have had an opportunity to flourish.

During most of the Phanerozoic, however, the ocean was fully oxygenated, resulting in high Mo and low Fe concentrations. Organisms with plastids derived from secondary red algal symbiosis, especially diatoms, are also well-adapted to cope with low Fe during NO_3^- assimilation in the modern ocean (Maldonado & Price, 1996; Maldonado & Price, 2000; Marchetti *et al.*, 2006). As shown in Fig. 7, the intracellular Fe content is lower in algae than cyanobacteria even when the expected Fe demand based on the N assimilation pathway being used is similar. This suggests that modern algae can subsist on lower Fe quotas than cyanobacteria, and may partially

explain the success of algae (especially diatoms) in the modern ocean. Metal-N limitation likely affected not only competition between eukaryotes versus prokaryotes but also competition between algal divisions (e.g. red versus green algae). The differences in metal concentrations between open ocean and coastal environments likely contributed to the shift from green to red algal dominance after the P/T boundary. Red algae, requiring higher concentrations of Mo, could have survived in oxic coastal environments (Quigg *et al.*, 2003), allowing their radiation in the Mesozoic and continued domination of the marine environment to the present day (Falkowski *et al.*, 2004). Their current prevalence may in part reflect their ability to utilize Cu to assimilate NH_4^+ from amino acids (see DON Assimilation section). Cu would have been scarce before the oceans became fully oxygenated (Saito *et al.*, 2003). Thus, in part the success of the modern phytoplankton population may reflect their adaptation to low Fe, and high Mo and Cu concentrations.

CONCLUSIONS AND FUTURE DIRECTIONS

By coupling geochemical, biochemical and genetic evidence, we have shown that the use of metals in N-assimilation metalloenzymes maps well to the emerging record of changing metal availability in the oceans. In the Archean, abundant Fe and small amounts of bioavailable NH_4^+ and NO_2^- led to the evolution of NH_4^+ and NO_2^- assimilation pathways in cyanobacteria (Fig. 5). These were insufficient to support the N demands of the growing microbial community, favoring the evolution of N_2 -fixing nitrogenase enzymes. Fe-Fe and Fe-V nitrogenases may have existed in ancient cyanobacteria, but a 'whiff' of O_2 (and Mo) at the Archean-Proterozoic boundary (2.5 Ga), and perhaps in earlier times as well, may have spurred the evolution of Mo-bearing nitrate reductase enzymes. Eventually, as Mo became available in the oceans, Fe-Fe and Fe-V nitrogenases were supplanted by the more efficient Mo-Fe version. It is, therefore, likely that most N assimilation proteins evolved by the time of the GOE. Thereafter, the intersection of metal availability with the differential metal requirements of different organisms may have contributed strongly to competition between prokaryotes versus eukaryotes. For example, cyanobacteria, which fared better than algae throughout the Proterozoic, are able to grow even at very low Mo concentrations on either N_2 or NO_3^- . This ability could have been a competitive advantage at a time when oceans were low in Mo compared to today (Scott *et al.*, 2008). If this ecological history is in fact related to the history of metal abundances then the rise of Mo in the Neoproterozoic could have contributed to the switch of the dominant phytoplankton community from cyanobacteria to green algae.

Future efforts should aim to more precisely determine the metal concentrations of ancient oceans, particularly before and after major evolutionary transitions, as well as the intracellular metal quotas and environmental metal abundances required to




Fig. 7 Comparison of Fe requirements for assimilation of different N sources for photosynthetic (A) prokaryotes (cyanobacteria) and (B) eukaryotes (secondary red algae). Bar graphs show the distribution of Fe atoms present in enzymes used to assimilate N_2 (prokaryotes only; Nif and GltS), NO_3^- (NarB in prokaryotes/NR in eukaryotes, NiR and GltS in prokaryotes/GltBS in eukaryotes), NO_2^- (NiR and GltS in prokaryotes/GltBD in eukaryotes), and NH_4^+ (GltS in prokaryotes/GltBD in eukaryotes). Note that the total Fe atoms used to assimilate inorganic N is greater in eukaryotes than prokaryotes for all N sources (aside from N_2) because of the 11 Fe atoms in GltBD, versus only 3 in GltS. Red data points represent intracellular Fe normalized to C for growth on different N sources at growth-limiting concentrations of Fe. Data sources: (a) N_2 -supported growth of cyanobacterium *Trichodesmium* IMS 101 (Kustka *et al.*, 2003b); NO_3^- -supported growth of cyanobacterium *Synechococcus* sp. (Kudo & Harrison, 1997); NH_4^+ -supported growth of species cyanobacterium *Trichodesmium* IMS 101 (Kustka *et al.*, 2003b) and *Synechococcus* sp. (Kudo & Harrison, 1997). (b) NO_3^- and NH_4^+ -supported growth of marine diatoms *Thalassiosira pseudonana* and *T. weissfloggi* (Maldonado & Price, 1996). (c) Intracellular Fe versus Fe atoms per N assimilation pathway for prokaryotes and eukaryotes. Note that intracellular Fe increases with greater numbers of Fe atoms per pathway, and that for a given number of Fe atoms in a pathway, prokaryotes have higher intracellular Fe quotas. This may reflect eukaryotic adaptation to the Phanerozoic ocean, where Fe levels are low, versus cyanobacterium adaptation to the Precambrian ocean, where Fe levels were higher.

support N_2 fixation and eukaryotic NO_3^- assimilation. Better understanding of metal concentrations through time, coupled with laboratory experiments to explore the physiological changes that organisms experience at representative metal concentrations, is needed to determine how biological pathways coevolved with changing Earth environments. Such an integrated geobiological approach has been quite informative in revealing the connections between changes in PO_2 , ocean sulfur abundance and microbial evolution through time (Canfield *et al.*, 2000; Shen *et al.*, 2001; Habicht *et al.*, 2002; Farquhar *et al.*, 2003; Johnston *et al.*, 2005). That precedent should serve as a model for further study of the evolutionary consequences of changing metal abundances.

In the succeeding discussion, we highlight examples of a larger set of themes that should drive future geochemical and

geobiological studies. We touch on several crucial future directions in geochemical research and then focus on geobiological investigations.

- Ocean Biogeochemical Evolution. We now have a crude but expanding picture of how marine Fe and Mo concentrations have changed through time (Fig. 1) (Anbar & Knoll, 2002; Rouxel *et al.*, 2005; Anbar *et al.*, 2007; Scott *et al.*, 2008), but little is known about the way that other metals have varied in the past. Emerging data indicate that Ni may have been higher in the Archean than in the Proterozoic and Phanerozoic oceans (Konhauser *et al.*, in press). Theoretical arguments suggest that Cu was lowest during the Proterozoic (Saito *et al.*, 2003; Zerkle *et al.*, 2005; Buick, 2007). Future geochemical studies and data compilations are needed to explore these and other such possibilities.

Further work is also required to gain greater knowledge of N geochemistry through time, where knowledge remains rudimentary. We do not currently have a clear picture of when certain metabolisms appeared (e.g. denitrification). Geochemical methods such as measurement of N isotopic compositions in sedimentary rocks have been useful in the past (Beaumont & Robert, 1999); more efforts are needed as analytical abilities advance. Previous studies have explored how marine inorganic N concentrations varied through ocean history (Beaumont & Robert, 1999; Papineau *et al.*, 2005). Much less is known about changes in DON (especially urea and amino acids) through time. This has important implications for metal requirements for N assimilation, as DON assimilation requires Ni and Cu, whereas DIN assimilation requires only Fe and Mo. Both modeling and geochemical studies are required to explore the marine speciation of N through time.

- **Microbial Metal Processing.** Better understanding of the strategies for uptake and regulation of metals that were scarce in ancient oceans can provide insight into the selection pressures exerted by metal scarcity on ancient ocean ecosystems, and consequent adaptations. We now know that uptake and regulation mechanisms can be diverse and complex. Today, marine photoautotrophs maximize their ability to acquire Fe by excreting siderophores and expressing high-affinity uptake systems to acquire Fe at minute quantities from seawater (Hutchins *et al.*, 1999; Maldonado & Price, 2001; Kraemer *et al.*, 2005). Marine diazotrophic cyanobacteria cycle Fe between nitrogenase and photosynthetic metalloproteins on a diurnal cycle, thus minimizing overall Fe requirements (Berman-Frank *et al.*, 2001; Tuit *et al.*, 2004; Saito *et al.*, 2008). Fe and N are also linked through regulatory networks that coordinate Fe uptake in accordance with requirements for N assimilatory enzymes (Maldonado & Price, 2000; López-Gomollón *et al.*, 2007). Additionally, some cyanobacteria store Fe in ferritin as a hedge against Fe scarcity (Castruita *et al.*, 2006, 2007). Other adaptations involve enzyme substitutions. For example, flavodoxin, an alternative to ferredoxin (see Nitrogen Fixation section) that does not incorporate Fe, is expressed by numerous phytoplankton when Fe concentrations are low (La Roche *et al.*, 1996).

Analogous intricate metal processing mechanisms could have evolved in ancient oceans starved of Mo, Cu, or other metals, and may have been inherited by modern cyanobacteria. This evolutionary legacy should be explored in order to understand how metals required for N assimilation (especially Mo, Cu, and Ni) are taken up and allocated intracellularly when scarce in the environment. For instance, a Mo storage protein has recently been discovered in a soil bacterium (Fenske *et al.*, 2005; Schemberg *et al.*, 2007). Such a protein may also be present even in the marine cyanobacterial diazotrophs, which possess higher Mo quotas than can be accounted for if nitrogenase is the sole Mo protein (Tuit *et al.*, 2004). Recent work has shown that molybdenum can be complexed by siderophores released by soil bacteria, allowing these organ-

isms to scavenge trace amounts of Mo (Bellenger *et al.*, 2008). Further research is required to understand the intracellular allocation of metals other than Fe in representative organisms from both modern and ancient environments.

- **Secondary Metal Requirements.** As N and metal metabolisms are explored with ever-increasing detail, utilization of metals for purposes secondary to direct N assimilation have been discovered. These metalloenzymes likely contribute to even higher metal requirements for N assimilation than previous recognized. For example, multi-Cu ferroxidases are used for oxidation of Fe(II) to Fe(III) in Fe-limited diatom cultures containing organically-bound Fe (Maldonado *et al.*, 2006). Thus, Cu is required in order for diatoms to access low levels of Fe required for acquisition of N in any form. Cu is also required for the biosynthesis of molybdopterin cofactors used in NR in higher plants (Kuper *et al.*, 2004). It is unknown whether cyanobacterial and algal molybdopterin biosynthesis also requires Cu. This represents an important future research direction, as an additional Cu requirement for NO_3^- assimilation would likely have had impacts on biological N acquisition in ancient oceans.

- **Metal Quotas.** The first comprehensive study of intracellular metal quotas in photoautotrophs provided an average trace element stoichiometry (Ho *et al.*, 2003). However, these stoichiometries can change under different nutrient regimes. This allows their use as a measure of trace element demand. As shown in Fig. 7, intracellular Fe quotas map well to expectations of Fe demand based on N-assimilation pathways, but show overall higher Fe demand for prokaryotes versus eukaryotes. Studies of intracellular Mo, Ni and Cu demand and partitioning are just beginning. Intracellular Mo has been shown to be higher in N_2 -fixing versus NO_3^- -grown cyanobacteria (Tuit *et al.*, 2004). Early studies showed that several species of algae had an absolute requirement for Mo when grown on NO_3^- as the sole N source (Walker, 1953; Loneragan & Arnon, 1954; Arnon *et al.*, 1955; Wallen & Cartier, 1975). Future studies could investigate the Mo requirements for NO_3^- assimilation for eukaryotes compared with those of cyanobacteria in order to test the hypothesis that eukaryotic acquisition of NO_3^- requires more Mo than prokaryotic assimilation. The Cu requirements for amine oxidation in algae could be measured in order to ascertain whether such low Cu concentrations could have limited N assimilation from amines in ancient oceans.

Finally, a fundamental question in geobiology is whether global environmental changes – like the rise of O_2 – trigger evolutionary innovation, or whether such changes enable pathways that evolved earlier, in unusual niches, to radiate (Raymond & Segre, 2006). Such distinctions remain unclear in the multi-billion-year history of N assimilation protein evolution, especially in the case of the different nitrate reductases. More precise phylogenetic dating of the evolution of these enzymes, coupled with more detailed geochemical knowledge of the Archean biosphere, will help elucidate this question.

ACKNOWLEDGEMENTS

Thanks to helpful discussions with John Berges, Robert Blankenship, Christopher Dupont, Tatsuo Omata, Jason Raymond, Luis Rubio, and William Sunda. Two anonymous reviewers provided insightful comments. Thanks to Susan Selkirk for the figure artwork. Gwyneth Gordon and Anthony Michaud aided with molybdenum and nitrogen concentration analyses. J.B.G. was supported on a National Science Foundation Graduate Student Research Fellowship (Geosciences-Geochemistry #2006038382), and F.W.-S. was supported by a National Science Foundation Minority Postdoctoral Fellowship (DBI-0511972). This work was also supported by grants from the Agouron Institute and the NASA Astrobiology Program to A.D.A.

REFERENCES

- Achterberg EP, van den Berg CMG, Boussemart M, Davison W (1997) Speciation and cycling of trace metals in Esthwaite Water: a productive English lake with seasonal deep-water anoxia. *Geochimica et Cosmochimica Acta* **61**, 5233–5253.
- Achterberg EP, Colombo C, van den Berg CMG (1999) The distribution of dissolved Cu, Zn, Ni, Co and Cr in English coastal surface waters. *Continental Shelf Research* **19**, 537–558.
- Ahmed J, Spiller H (1976) Purification and some properties of the nitrate reductase from *Ankistrodesmus braunii*. *Plant and Cell Physiology* **17**, 1–10.
- Algeo TJ (2004) Can marine anoxic events draw down the trace element inventory of seawater? *Geology* **32**, 1057–1060.
- Allen AE, Ward BB, Song B (2005) Characterization of diatom (*Bacillariophyceae*) nitrate reductase genes and their detection in marine phytoplankton communities. *Journal of Phycology* **41**, 95–104.
- Amy NK, Garrett RH (1974) Purification and characterization of the nitrate reductase from the diatom *Thalassiosira pseudonana*. *Plant Physiology* **54**, 629–637.
- Anbar AD (2008) Elements and evolution. *Science* **322**, 1481–1483.
- Anbar AD, Gordon GW (2008) Redox renaissance. *Geology* **36**, 271.
- Anbar AD, Knoll AH (2002) Proterozoic ocean chemistry and evolution: a bioinorganic bridge? *Science* **297**, 1137–1142.
- Anbar AD, Rouxel O (2007) Metal stable isotopes in paleoceanography. *Annual Reviews of Earth and Planetary Sciences* **35**, 717–746.
- Anbar AD, Duan Y, Lyons TW, Arnold GL, Kendall B, Creaser RA, Kaufman AJ, Gordon GW, Scott C, Garvin J (2007) A whiff of oxygen before the great oxidation event? *Science* **317**, 1903–1906.
- Andreini C, Banci L, Bertini I, Rosato A (2008) Occurrence of copper proteins through the three domains of life: a bioinformatic approach. *Journal of Proteome Research* **7**, 209–216.
- Antia NJ, Harrison PJ, Oliveira L (1991) The role of dissolved organic nitrogen in phytoplankton nutrition, cell biology and ecology. *Phycologia* **30**, 1–89.
- Antipov AN, Sorokin DY, L'Vov NP, Kuenen JG (2003) New enzyme belonging to the family of molybdenum-free nitrate reductases. *Biochemical Journal* **369**, 185–189.
- Antipov AN, Morozkina EV, Sorokin DY, Golubeva LI, Zvyagilskaya RA, L'Vov NP (2005) Characterization of molybdenum-free nitrate reductase from haloalkalophilic bacterium *Halomonas* sp strain AGJ 1-3. *Biochemistry-Moscow* **70**, 799–803.
- Antipova AN, Lyalikova NN, Khijniak TV, L'Vov NP (1998) Molybdenum-free nitrate reductases from vanadate-reducing bacteria. *FEBS Letters* **441**, 257–260.
- Aparicio PJ, Cardenas J, Zumft WG, Vega JM, Herrera J, Paneque A, Losada M (1971) Molybdenum and iron as constituents of the enzymes of the nitrate reducing system from *Chlorella*. *Phytochemistry* **10**, 1487–1495.
- Argall ME, Smith GD, Stamford NPJ, Youens BN (1992) Purification and properties of urease from the cyanobacterium *Anabaena cylindrica*. *Biochemistry International* **27**, 1027–1036.
- Arizmendi JM, Serra JL (1990) Purification and some properties of the nitrite reductase from the cyanobacterium *Phormidium laminosum*. *Biochimica et Biophysica Acta* **2**, 237–244.
- Arnold GL, Anbar AD, Barling J, Lyons TW (2004) Molybdenum isotope evidence for widespread anoxia in mid-Proterozoic oceans. *Science* **304**, 87–90.
- Arnon DI, Ichioka PS, Wessel G, Fujiwara A, Wolley JT (1955) Molybdenum in relation to nitrogen metabolism I. Assimilation of nitrate nitrogen by *Scenedesmus*. *Physiologia Plantarum* **8**, 538.
- Arrigo KR (2005) Marine microorganisms and global nutrient cycles. *Nature* **437**, 349–355.
- Axler RP, Reuter JE (1996) Nitrate uptake by phytoplankton and periphyton: whole-lake enrichments and mesocosm ¹⁵N experiments in an oligotrophic lake. *Limnology and Oceanography* **41**, 659–671.
- Axler RP, Gersberg RM, Goldman CR (1980) Stimulation of nitrate uptake and photosynthesis by molybdenum in Castle Lake, California. *Canadian Journal of Fisheries and Aquatic Sciences* **37**, 707–712.
- Bachmann RW, Goldman CR (1964) The determination of microgram quantities of molybdenum in natural waters. *Limnology and Oceanography* **9**, 143–146.
- Baumont V, Robert F (1999) Nitrogen isotope ratios of kerogens in Precambrian cherts: a record of the evolution of atmosphere chemistry? *Precambrian Research* **96**, 63–82.
- Bellenger JP, Wichard T, Kustka AB, Kraepiel AM (2008) Uptake of molybdenum and vanadium by a nitrogen-fixing soil bacterium using siderophores. *Nature Geoscience* **1**, 243–246.
- Berges JA (1997) Algal nitrate reductases. *European Journal of Phycology* **32**, 3–8.
- Berges JA, Mulholland M (2008) Enzymes and N cycling. In *Nitrogen in the Marine Environment*, second edition (eds Capone DG, Bronk DA, Mulholland M, Carpenter EJ). Academic Press, New York, pp. 1361–1420.
- Berman-Frank I, Cullen JT, Shaked Y, Sherrell RM, Falkowski PG (2001) Iron availability, cellular iron quotas, and nitrogen fixation in *Trichodesmium*. *Limnology and Oceanography* **46**, 1249–1260.
- Berman-Frank I, Lundgren P, Falkowski P (2003) Nitrogen fixation and photosynthetic oxygen evolution in cyanobacteria. *Research in Microbiology* **154**, 157–164.
- Berman-Frank I, Quigg A, Finkel ZV, Irwin AJ, Haramaty L (2007) Nitrogen-fixation strategies and Fe requirements in cyanobacteria. *Limnology and Oceanography* **52**, 2260–2269.
- Beudeker RF, Tabita FR (1985) Characterization of glutamine synthetase isoforms from *Chlorella*. *Plant Physiology* **77**, 791–794.
- Binda C, Bossi RT, Wakatsuki S, Arzt S, Coda A, Curti B, Vanoni MA, Mattevi A (2000) Cross-talk and ammonia channeling between active centers in the unexpected domain arrangement of glutamate synthase. *Structure* **8**, 1299–1308.

- Bishop PE, Jarlenski DM, Hetherington DR (1980) Evidence for an alternative nitrogen fixation system in *Azotobacter vinelandii*. *Proceedings of the National Academy of Sciences, USA* **77**, 7342.
- Blanco F, Alana A, Llama MJ, Serra JL (1989) Purification and properties of glutamine synthetase from the non-N₂-fixing cyanobacterium *Phormidium laminosum*. *Journal of Bacteriology* **171**, 1158–1165.
- Blankenship RE (1992) Origin and early evolution of photosynthesis. *Photosynthesis Research* **33**, 91–111.
- Blankenship RE (2001) Molecular evidence for the evolution of photosynthesis. *Trends in Plant Science* **6**, 4–6.
- Boison G, Steingen C, Stal LJ, Bothe H (2006) The rice field cyanobacteria *Anabaena azotica* and *Anabaena* sp. CH1 express vanadium-dependent nitrogenase. *Archives of Microbiology* **186**, 367–376.
- Bortels H (1930) Molybdan als Katalysator bei der biologischen Stickstoffbindung. *Archives of Microbiology* **1**, 333–342.
- Bortels H (1940) Über die Bedeutung des Molybdäns für stickstoffbindende Nostocaceen. *Archives of Microbiology* **11**, 155–186.
- Boyd SR, Philippot P (1998) Precambrian ammonium biogeochemistry: a study of the Moine metasediments, Scotland. *Chemical Geology* **144**, 257–268.
- Brocks JJ, Logan GA, Buick R, Summons RE (1999) Archean molecular fossils and the early rise of eukaryotes. *Science* **285**, 1033–1036.
- Brocks JJ, Buick R, Logan GA, Summons RE (2003a) Composition and syngeneity of molecular fossils from the 2.78 to 2.45 billion-year-old Mount Bruce Supergroup, Pilbara Craton, Western Australia. *Geochimica et Cosmochimica Acta* **67**, 4289–4319.
- Brocks JJ, Buick R, Summons RE, Logan GA (2003b) A reconstruction of Archean biological diversity based on molecular fossils from the 2.78 to 2.45 billion-year-old Mount Bruce Supergroup, Hamersley Basin, Western Australia. *Geochimica et Cosmochimica Acta* **67**, 4321–4335.
- Brocks JJ, Love GD, Summons RE, Knoll AH, Logan GA, Bowden SA (2005) Biomarker evidence for green and purple sulphur bacteria in a stratified Palaeoproterozoic sea. *Nature* **437**, 866–870.
- Bronk DA, See JH, Bradley P, Killberg L (2007) DON as a source of bioavailable nitrogen for phytoplankton. *Biogeosciences* **4**, 283–296.
- Bruland KW (1980) Oceanographic distributions of cadmium, zinc, nickel, and copper in the North Pacific. *Earth and Planetary Science Letters* **47**, 176–198.
- Bruland KW, Rue EL, Smith GJ (2001) Iron and macronutrients in California coastal upwelling regimes: implications for diatom blooms. *Limnology and Oceanography* **46**, 1661–1674.
- Buick R (1992) The antiquity of oxygenic photosynthesis: evidence from stromatolites in sulphate-deficient Archean lakes. *Science* **255**, 74–77.
- Buick R (2007) Did the Proterozoic Canfield Ocean cause a laughing gas greenhouse? *Geobiology* **5**, 97–100.
- Buick R (2008) When did oxygenic photosynthesis evolve? *Philosophical Transactions of the Royal Society B: Biological Sciences* **363**, 2731–2743.
- Burke DH, Hearst JE, Sidow A (1993) Early evolution of photosynthesis: clues from nitrogenase and chlorophyll iron proteins. *Proceedings of the National Academy of Sciences, USA* **90**, 7134.
- Burns RC, Fuchsman WH, Hardy RW (1971) Nitrogenase from vanadium-grown *Azotobacter*: isolation, characteristics, and mechanistic implications. *Biochemical and Biophysical Research Communications* **42**, 353–358.
- Cameron EM, Garrels RM (1980) Geochemical compositions of some Precambrian shales from the Canadian Shield. *Chemical Geology* **28**, 181–197.
- Campbell WH (1999) Nitrate reductase structure, function and regulation: bridging the gap between biochemistry and physiology. *Annual Review of Plant Physiology and Plant Molecular Biology* **50**, 277–303.
- Canfield DE (1998) A new model for Proterozoic ocean chemistry. *Nature* **396**, 450–453.
- Canfield DE, Habicht KS, Thamdrup B (2000) The Archean sulfur cycle and the early history of atmospheric oxygen. *Science* **288**, 658–661.
- Canfield DE, Poulton SW, Knoll AH, Narbonne GM, Ross G, Goldberg T, Strauss H (2008) Ferruginous conditions dominated later Neoproterozoic deep-water chemistry. *Science* **321**, 949–952.
- Capone DG, Carpenter EJ (1982) Nitrogen fixation in the marine environment. *Science* **217**, 1140–1142.
- Capone DG, Zehr JP, Paerl HW, Bergman B, Carpenter EJ (1997) *Trichodesmium*, a globally significant marine cyanobacterium. *Science* **276**, 1221–1229.
- Cárdenas J, Rivas J, Paneque A, Losada M (1971) Molybdenum and the nitrate-reducing system from *Chlorella*. *Archives of Microbiology* **79**, 367–376.
- Cárdenas J, Rivas J, Paneque A, Losada M (1972) Effect of iron supply on the activities of the nitrate-reducing system from *Chlorella*. *Archives of Microbiology* **81**, 260–263.
- Carpenter EJ, Bergman B, Dawson R, Siddiqui PJ, Soderback E, Capone DG (1992) Glutamine synthetase and nitrogen cycling in colonies of the marine diazotrophic cyanobacteria *Trichodesmium* spp. *Applied and Environmental Microbiology* **58**, 3122–3129.
- Carvajal N, Fernandez M, Rodriguez JP, Donoso M (1982) Urease of *Spirulina maxima*. *Phytochemistry* **21**, 2821–2823.
- Castruita M, Saito M, Schottel PC, Elmegeen LA, Myneni S, Stiefel EI, Morel FMM (2006) Overexpression and characterization of an iron storage and DNA-binding Dps protein from *Trichodesmium erythraeum*. *Applied and Environmental Microbiology* **72**, 2918–2924.
- Castruita M, Elmegeen LA, Shaked Y, Stiefel EI, Morel FMM (2007) Comparison of the kinetics of iron release from a marine (*Trichodesmium erythraeum*) Dps protein and mammalian ferritin in the presence and absence of ligands. *Journal of Inorganic Biochemistry* **101**, 1686–1691.
- Catling DC, Claire MW (2005) How Earth's atmosphere evolved to anoxic state: a status report. *Earth and Planetary Science Letters* **237**, 1–20.
- Clayton JR, Ahmed SI (1986) Detection of glutamate synthase (GOGAT) activity in phytoplankton: evaluation of cofactors and assay optimization. *Marine Ecology Progress Series* **32**, 115–122.
- Chisnell JR, Premakumar R, Bishop PE (1988) Purification of a second alternative nitrogenase from a *nifHDK* deletion strain of *Azotobacter vinelandii*. *Journal of Bacteriology* **170**, 27–33.
- Coale KH, Johnson KS, Fitzwater SE, Gordon RM, Tanner S, Chavez FP, Ferioli L, Sakamoto C, Rogers P, Millero F (1996) A massive phytoplankton bloom induced by an ecosystem-scale iron fertilization experiment in the equatorial Pacific Ocean. *Nature* **383**, 495–501.
- Cole JJ, Lane JM, Marino R, Howarth RW (1993) Molybdenum assimilation by cyanobacteria and phytoplankton in freshwater and salt water. *Limnology and Oceanography* **38**, 25–35.
- Collier RW (1985) Molybdenum in the northeast Pacific Ocean. *Limnology and Oceanography* **30**, 1351–1354.
- Collier JL, Brahamsha B, Palenik B (1999) The marine cyanobacterium *Synechococcus* sp. WH7805 requires urease

- (urea amidohydrolase, EC 3.5.1.5) to utilize urea as a nitrogen source: molecular-genetic and biochemical analysis of the enzyme. *Microbiology* **145**, 447–459.
- Cona A, Rea G, Angelini R, Federico R, Tavloraki P (2006) Functions of amine oxidases in plant development and defence. *Trends in Plant Science* **11**, 80–88.
- Cullimore JV, Sims AP (1981) Occurrence of two forms of glutamate synthase in *Chlamydomonas reinhardtii*. *Phytochemistry* **20**, 597–600.
- Davy R (1983) A geochemical study of the Mount McRae shale and the upper part of the Mount Sylvania Formation in Core RD1, Rhodes Ridge, Western Australia. *Geological Survey of Western Australia Record* **1983/3**.
- Dellwig O, Beck M, Lemke A, Lunau M, Kolditz K, Schnetger B, Brumsack HJ (2007) Non-conservative behaviour of molybdenum in coastal waters: coupling geochemical, biological, and sedimentological processes. *Geochimica et Cosmochimica Acta* **71**, 2745–2761.
- Dias JM, Than ME, Humm A, Huber R, Bourenkov GP, Bartunik HD, Bursakov S, Calvete J, Caldeira J, Carneiro C, Moura JGG, Moura I, Romao M (1999) Crystal structure of the first dissimilatory nitrate reductase at 1.9 Å solved by MAD methods. *Structure* **7**, 65–79.
- Dupont CL, Barbeau K, Palenik B (2007) Ni uptake and limitation in marine *Synechococcus*. *Applied and Environmental Microbiology* **74**, 23–31.
- Dyhrman ST, Anderson DM (2003) Urease activity in cultures and field populations of the toxic dinoflagellate *Alexandrium*. *Limnology and Oceanography* **48**, 647–655.
- Eady RR (1996) Structure-function relationships of alternative nitrogenases. *Chemical Reviews* **96**, 3013–3030.
- Egleston ES, Morel FMM (2008) Nickel limitation and zinc toxicity in a urea-grown diatom. *Limnology and Oceanography* **53**, 2462–2471.
- Eigenbrode JL, Freeman KH (2006) Late Archean rise of aerobic microbial ecosystems. *Proceedings of the National Academy of Sciences, USA* **103**, 15759–15764.
- El Alaoui S, Diez J, Toribio F, Gomez-Baena G, Dufresne A, Garcia-Fernandez JM (2003) Glutamine synthetase from the marine cyanobacteria *Prochlorococcus* spp. characterization, phylogeny and response to nutrient limitation. *Environmental Microbiology* **5**, 412–423.
- Falkowski PG (1983) Enzymology of nitrogen assimilation. In *Nitrogen in the Marine Environment* (eds Carpenter EJ, Capone DG). Academic Press, New York, pp. 839–868.
- Falkowski PG (1997) Evolution of the nitrogen cycle and its influence on the biological sequestration of CO₂ in the ocean. *Nature* **387**, 272–275.
- Falkowski PG, Godfrey LV (2008) Electrons, life and the evolution of the Earth's oxygen cycle. *Philosophical Transactions of the Royal Society of London Series B, Biological Sciences* **363**, 2705–2716.
- Falkowski PG, Katz ME, Knoll AH, Quigg A, Raven JA, Schofield O, Taylor FJR (2004) The evolution of modern eukaryotic phytoplankton. *Science* **305**, 354–360.
- Fani R, Gallo R, Liò P (2000) Molecular evolution of nitrogen fixation: the evolutionary history of the *nifD*, *nifK*, *nifE*, and *nifN* genes. *Journal of Molecular Evolution* **51**, 1–11.
- Farquhar J, Johnston DT, Wing BA, Habicht KS, Canfield DE, Airieau S, Thieme MH (2003) Multiple sulphur isotopic interpretations of biosynthetic pathways: implications for biological signatures in the sulphur isotope record. *Geobiology* **1**, 27–36.
- Feng DF, Cho G, Doolittle RF (1997) Determining divergence times with a protein clock: update and reevaluation. *Proceedings of the National Academy of Sciences, USA* **94**, 13028–13033.
- Fennel K, Follows M, Falkowski PG (2005) The co-evolution of the nitrogen, carbon and oxygen cycles in the Proterozoic ocean. *American Journal of Science* **305**, 526.
- Fenske D, Gnida M, Schneider K, Meyer-Klaucke W, Schemberg J, Henschel V, Meyer AK, Knochel A, Müller A (2005) A new type of metalloprotein: the Mo storage protein from *Azotobacter vinelandii* contains a polynuclear molybdenum–oxide cluster. *ChemBioChem* **6**, 405–413.
- Fike DA, Grotzinger JP, Pratt LM, Summons RE (2006) Oxidation of the Ediacaran ocean. *Nature* **444**, 744–747.
- Fischer K, Barbier GG, Hecht HJ, Mendel RR, Campbell WH, Schwarz G (2005) Structural basis of eukaryotic nitrate reduction: crystal structures of the nitrate reductase active site. *The Plant Cell Online* **17**, 1167–1179.
- Florencio FJ, Ramos JL (1985) Purification and characterization of glutamine synthetase from the unicellular cyanobacterium *Anacystis nidulans*. *Biochimica et Biophysica Acta* **838**, 39–48.
- Garcia-Dominguez M, Reyes JC, Florencio FJ (1997) Purification and characterization of a new type of glutamine synthetase from cyanobacteria. *European Journal of Biochemistry* **244**, 258–264.
- Garvin J, Buick R, Anbar AD, Arnold GL, Kaufman AJ (2009) Isotopic evidence for an aerobic nitrogen cycle in the latest Archean. *Science* **323**, 1045–1048.
- Georgiadis MM, Komiya H, Chakrabarti P, Woo D, Kornuc JJ, Rees DC (1992) Crystallographic structure of the nitrogenase iron protein from *Azotobacter vinelandii*. *Science* **257**, 1653–1659.
- Glibert PM, Harrison J, Heil C, Seitzinger S (2006) Escalating worldwide use of urea – a global change contributing to coastal eutrophication. *Biogeochemistry* **77**, 441–463.
- Goldman CR (1960) Molybdenum as a factor limiting primary productivity in Castle Lake, California. *Science* **132**, 1016–1017.
- Gollan U, Schneider K, Muller A, Schuddekopf K, Klipp W (1993) Detection of the *in vivo* incorporation of a metal cluster into a protein. The FeMo cofactor is inserted into the FeFe protein of the alternative nitrogenase of *Rhodospirillum rubrum*. *European Journal of Biochemistry* **215**, 25–35.
- Gotor C, Martinez-Rivas JM, Marquez AJ, Vega JM (1990) Functional properties of purified ferredoxin–glutamate synthase from *Chlamydomonas reinhardtii*. *Phytochemistry* **29**, 711–717.
- Grice K, Cao C, Love GD, Bottcher ME, Twitchett RJ, Grosjean E, Summons RE, Turgeon SC, Dunning W, Jin Y (2005) Photic zone euxinia during the Permian–Triassic superanoxic event. *Science* **307**, 706–709.
- Grzebyk D, Schofield O, Vetrani C, Falkowski PG (2003) The Mesozoic radiation of eukaryotic algae: the portable plastid hypothesis. *Journal of Phycology* **39**, 259–267.
- Guerrero MG, Vega JM, Losada M (1981) The assimilatory nitrate-reducing system and its regulation. *Annual Review of Plant Physiology and Plant Molecular Biology* **32**, 169–204.
- Habicht KS, Gade M, Thamdrup B, Berg P, Canfield DE (2002) Calibration of sulfate levels in the Archean ocean. *Science* **298**, 2372.
- Hallenbeck PC, Kostel PJ, Benemann JR (1979) Purification and properties of nitrogenase from the cyanobacterium *Anabaena cylindrica*. *European Journal of Biochemistry* **98**, 275–284.
- Hattori A, Myers J (1967) Reduction of nitrate and nitrite by subcellular preparations of *Anabaena cylindrica*. 2. Reduction of nitrate to nitrite. *Plant and Cell Physiology* **8**, 327–337.
- Hedges SB, Chen H, Kumar S, Wang DYC, Thompson AS, Watanabe H (2001) A genomic timescale for the origin of eukaryotes. *BMC Evolutionary Biology* **1**.
- van den Heuvel RHH, Ferrari D, Bossi RT, Ravasio S, Curti B, Vanoni MA, Florencio FJ, Mattevi A (2002) Structural studies on the synchronization of catalytic centers in glutamate synthase. *Journal of Biological Chemistry* **277**, 24579–24583.

- van den Heuvel RHH, Svergun DI, Petoukhov MV, Coda A, Curti B, Ravasio S, Vanoni MA, Mattevi A (2003) The active conformation of glutamate synthase and its binding to ferredoxin. *Journal of Molecular Biology* **330**, 113–128.
- Hirasawa M, Rubio LM, Griffin JL, Flores E, Herrero A, Li J, Kim SK, Hurley JK, Tollin G, Knaff DB (2004) Complex formation between ferredoxin and *Synechococcus* ferredoxin: nitrate oxidoreductase. *Biochimica et Biophysica Acta* **1608**, 155–162.
- Ho TY, Quigg A, Finkel ZV, Milligan AJ, Wyman K, Falkowski PG, Morel FMM (2003) The elemental composition of some marine phytoplankton. *Journal of Phycology* **39**, 1145–1159.
- Holland HD (1973) The oceans: a possible source of iron in iron-formations. *Economic Geology* **68**, 1169–1172.
- Holland HD (2002) Volcanic gases, black smokers, and the great oxidation event. *Geochimica et Cosmochimica Acta* **66**, 3811–3826.
- Holland HD (2006) The oxygenation of the atmosphere and oceans. *Philosophical Transactions of the Royal Society of London Series B, Biological Sciences* **361**, 903–915.
- Holm L, Sander C (1997) An evolutionary treasure: unification of a broad set of amidohydrolases related to urease. *Proteins* **28**, 72–82.
- Howard JB, Rees DC (1996) Structural basis of biological nitrogen fixation. *Chemical Reviews* **96**, 2965–2982.
- Howarth RW, Cole JJ (1985) Molybdenum availability, nitrogen limitation, and phytoplankton growth in natural waters. *Science* **229**, 653–655.
- Hutchins DA, Witter AE, Butler A, Luther GW III (1999) Competition among marine phytoplankton for different chelated iron species. *Nature* **400**, 858–861.
- Ida S, Mikami B (1983) Purification and characterization of assimilatory nitrate reductase from the cyanobacterium *Plectonema boryanum*. *Plant and Cell Physiology* **24**, 649–658.
- Jabri E, Carr MB, Hausinger RP, Karplus PA (1995) The crystal structure of urease from *Klebsiella aerogenes*. *Science* **268**, 998–1004.
- Jahns T, Schafer U, Kaltwasser H (1995) Heat-stable ureases from two filamentous cyanobacteria. *Microbiology* **141**, 737–741.
- Jepson BJN, Anderson LJ, Rubio LM, Taylor CJ, Butler CS, Flores E, Herrero A, Butt JN, Richardson DJ (2004) Tuning a nitrate reductase for function – the first spectropotentiometric characterization of a bacterial assimilatory nitrate reductase reveals novel redox properties. *Journal of Biological Chemistry* **279**, 32212–32218.
- Joerger RD, Bishop PE (1988) Bacterial alternative nitrogen fixation systems. *Critical Reviews in Microbiology* **16**, 1–14.
- Johnson KS, Gordon RM, Coale KH (1997) What controls dissolved iron concentrations in the world ocean? *Marine Chemistry* **57**, 137–161.
- Johnson CM, Beard BL, Roden EE (2008) The iron isotope fingerprints of redox and biogeochemical cycling in the modern and ancient Earth. *Annual Reviews of Earth and Planetary Sciences* **36**, 457–493.
- Johnston DT, Wing BA, Farquhar J, Kaufman AJ, Strauss H, Lyons TW, Kah LC, Canfield DE (2005) Active microbial sulfur disproportionation in the Mesoproterozoic. *Science* **310**, 1477–1479.
- Junium CK, Arthur MA (2007) Nitrogen cycling during the Cretaceous, Cenomanian–Turonian Oceanic Anoxic Event II. *Geochemistry Geophysics Geosystems* **8**, Q03002, doi: 10.1029/2006GC001328.
- Kashiyama Y, Ogawa NO, Kuroda J, Shiro M, Nomoto RT, Kitazato H, Ohkouchi N (2008) Diazotrophic cyanobacteria as the major photoautotrophs during mid-Cretaceous oceanic anoxic events: Nitrogen and carbon isotopic evidence from sedimentary porphyrin. *Organic geochemistry* **39**, 532–549.
- Kasting JF, Catling D (2003) Evolution of a habitable planet. *Annual Reviews in Astronomy and Astrophysics* **41**, 429–463.
- Kaufman AJ, Johnston DT, Farquhar J, Masterson AL, Lyons TW, Bates S, Anbar AD, Arnold GL, Garvin J, Buick R (2007) Late Archean biospheric oxygenation and atmospheric evolution. *Science* **317**, 1900–1903.
- Kentemich T, Danneberg G, Hundeshagen B, Bothe H (1988) Evidence for the occurrence of the alternative, vanadium-containing nitrogenase in the cyanobacterium *Anabaena variabilis*. *FEMS Microbiology Letters* **51**, 19–24.
- Kim J, Rees DC (1992) Crystallographic structure and functional implications of the nitrogenase molybdenum-iron protein from *Azotobacter vinelandii*. *Nature* **360**, 553–560.
- Kirschvink JL, Kopp RE (2008) Paleoproterozoic icehouses and the evolution of oxygen mediating enzymes: the case for a late origin of photosystem II. *Philosophical Transactions of the Royal Society of London Series B, Biological Sciences* **363**, 2755–2765.
- Kirschvink JL, Gaidos EJ, Bertani LE, Beukes NJ, Gutzmer J, Maepa LN, Steinberger RE (2000) Paleoproterozoic snowball Earth: extreme climatic and geochemical global change and its biological consequences. *Proceedings of the National Academy of Sciences, USA* **97**, 1400.
- Knoll AH (2007) The geological succession of primary producers in the oceans. In *Evolution of Primary Producers in the Sea* (eds Falkowski PG, Knoll AH). Elsevier Academic Press, London, UK, pp. 133–163.
- Konhauser KO, Pecoits E, Lalonde SV, Papineau D, Nisbet EG, Barley ME, Arndt NT, Zahnle K, Balz S. Oceanic nickel depletion and a methanogen famine. *Nature*, in press.
- Koper TE, El-Sheikh AF, Norton JM, Klotz MG (2004) Urease-encoding genes in ammonia-oxidizing bacteria. *Applied and Environmental Microbiology* **70**, 2342.
- Kraemer SM, Butler A, Borer P, Cervini-Silva J (2005) Siderophores and the dissolution of iron-bearing minerals in marine systems. *Reviews in Mineralogy and Geochemistry* **59**, 53–84.
- Kudo I, Harrison PJ (1997) Effect of iron nutrition of the marine cyanobacterium *Synechococcus* grown on different N sources and irradiances. *Journal of Phycology* **33**, 232–240.
- Kumar V, Dooley DM, Freeman HC, Guss JM, Harvey I, McGuirl MA, Wilce MCJ, Zubak VM (1996) Crystal structure of a eukaryotic (pea seedling) copper-containing amine oxidase at 2.2 Å resolution. *Structure* **4**, 943–955.
- Kump L (2005) Ironing out biosphere oxidation. *Science* **307**, 1058–1059.
- Kump LR, Barley ME (2007) Increased subaerial volcanism and the rise of atmospheric oxygen 2.5 billion years ago. *Nature* **448**, 1033–1036.
- Kump LR, Pavlov A, Arthur MA (2005) Massive release of hydrogen sulfide to the surface ocean and atmosphere during intervals of oceanic anoxia. *Geology* **33**, 397–400.
- Kuper J, Llamas A, Hecht HJ, Mendel RR, Schwarz G (2004) Structure of the molybdopterin-bound Cnx 1 G domain links molybdenum and copper metabolism. *Nature* **430**, 803–806.
- Kustka A, Sanudo-Wilhelmy S, Carpenter EJ, Capone DG, Raven JA (2003a) A revised estimate of the iron use efficiency of nitrogen fixation, with special reference to the marine cyanobacterium *Trichodesmium* spp. (Cyanophyta). *Journal of Phycology* **39**, 12–25.
- Kustka AB, Sanudo-Wilhelmy SA, Carpenter EJ, Capone D, Burns J, Sunda WG (2003b) Iron requirements for dinitrogen- and ammonium-supported growth in cultures of *Trichodesmium* (IMS 101): Comparison with nitrogen fixation rates and iron: carbon ratios of field populations. *Limnology and Oceanography* **48**, 1869–1884.

- Kuypers MMM, van Breugel Y, Schouten S, Erba E, Damste JSS (2004) N₂-fixing cyanobacteria supplied nutrient N for Cretaceous oceanic anoxic events. *Geology* **32**, 853–856.
- Kuznetsova S, Knaff DB, Hirasawa M, Lagoutte B, Setif P (2004) Mechanism of spinach chloroplast ferredoxin-dependent nitrite reductase: spectroscopic evidence for intermediate states. *Biochemistry* **43**, 510–517.
- La Roche J, Boyd PW, McKay RML, Geider RJ (1996) Flavodoxin as an in situ marker for iron stress in phytoplankton. *Nature* **382**, 802–805.
- Lomas MW, Lipschultz F (2006) Forming the primary nitrite maximum: nitrifiers or phytoplankton? *Limnology and Oceanography* **51**, 2453–2467.
- Loneragan J, Arnon D (1954) Molybdenum in the growth and metabolism of *Chlorella*. *Nature* **174**, 459.
- López-Gomollón S, Hernández JA, Pellicer S, Angarica VE, Peleato ML, Fillat MF (2007) Cross-talk between iron and nitrogen regulatory networks in *Anabaena* (*Nostoc*) sp. PCC 7120: Identification of overlapping genes in FurA and NtcA regulons. *Journal of Molecular Biology* **374**, 267–281.
- Lopez-Ruiz A, Verbelen JP, Bocanegra JA, Diez J (1991) Immunocytochemical localization of nitrite reductase in green algae. *Plant Physiology* **96**, 699–704.
- Lu G, Lindqvist Y, Schneider G, Dwivedi U, Campbell W (1995) Structural studies on corn nitrate reductase: refined structure of the cytochrome *b* reductase fragment at 2.5 Å, its ADP complex and an active-site mutant and modeling of the cytochrome *b* domain. *Journal of Molecular Biology* **248**, 931–948.
- Luque I, Flores E, Herrero A (1993) Nitrite reductase gene from *Synechococcus* sp. PCC 7942 – homology between cyanobacterial and higher-plant nitrite reductases. *Plant Molecular Biology* **21**, 1201–1205.
- Mackerras AH, Smith GD (1986) Urease activity of the cyanobacterium *Anabaena cylindrica*. *Journal of General Microbiology* **132**, 2749–2752.
- Madigan MT (1995) Microbiology of nitrogen fixation by anoxygenic photosynthetic bacteria. In *Anoxygenic Photosynthetic Bacteria* (eds Blankenship RE, Madigan MT, Bauer CE). Springer, The Netherlands, pp. 915–928.
- Maitz GS, Haas EM, Castric PA (1982) Purification and properties of the allophanate hydrolase from *Chlamydomonas reinhardtii*. *Biochimica et Biophysica Acta* **714**, 486–491.
- Maldonado MT, Price NM (1996) Influence of N substrate on Fe requirements of marine centric diatoms. *Marine Ecology Progress Series* **141**, 161–172.
- Maldonado MT, Price NM (2000) Nitrate regulation of Fe reduction and transport by Fe-limited *Thalassiosira oceanica*. *Limnology and Oceanography* **45**, 814–826.
- Maldonado MT, Price NM (2001) Reduction and transport of organically bound iron by *Thalassiosira oceanica* (Bacillariophyceae). *Journal of Phycology* **37**, 298–310.
- Maldonado MT, Allen AE, Chong JS, Lin K, Leus D, Karpenko N, Harris SL (2006) Copper-dependent iron transport in coastal and oceanic diatoms. *Limnology and Oceanography* **51**, 1729–1743.
- Mancinelli R, McKay C (1988) The evolution of nitrogen cycling. *Origins of Life and Evolution of the Biosphere* **18**, 311–325.
- Manzano C, Candau P, Gomezmoreno C, Relimpio AM, Losada M (1976) Ferredoxin-dependent photosynthetic reduction of nitrate and nitrite by particles of *Anacystis nidulans*. *Molecular and Cellular Biochemistry* **10**, 161–169.
- Marchetti A, Maldonado MT, Lane ES, Harrison PJ (2006) Iron requirements of the pennate diatom *Pseudo-nitzschia*: comparison of oceanic (high-nitrate, low-chlorophyll waters) and coastal species. *Limnology and Oceanography* **51**, 2092–2101.
- Marques S, Florencio FJ, Candau P (1992) Purification and characterization of the ferredoxin-glutamate synthase from the unicellular cyanobacterium *Synechococcus* sp PCC 6301. *European Journal of Biochemistry* **206**, 69–77.
- Marquez AJ, Galvan F, Vega JM (1984) Purification and characterization of the NADH-glutamate synthase from *Chlamydomonas reinhardtii*. *Plant Science Letters* **34**, 305–314.
- Martin JH, Fitzwater SE (1988) Iron deficiency limits phytoplankton growth in the north-east Pacific subarctic. *Nature* **331**, 341–343.
- Masepohl B, Drepper T, Klipp W (2004) Nitrogen fixation in the photosynthetic purple bacterium *Rhodobacter capsulatus*. In *Genetics and Regulation of Nitrogen Fixation in Free-Living Bacteria* (eds Klipp W, Masepohl B, Gallon JR, Newton WE). Springer, The Netherlands, pp. 141–173.
- Mather TA, Pyle DM, Allen AG (2004) Volcanic source for fixed nitrogen in the early Earth's atmosphere. *Geology* **32**, 905–908.
- McFadden KA, Huang J, Chu X, Jiang G, Kaufman AJ, Zhou C, Yuan X, Xiao S (2008) Pulsed oxidation and biological evolution in the Ediacaran Doushantuo Formation. *Proceedings of the National Academy of Sciences, USA* **105**, 3197.
- Meeks JC, Wolk CP, Lockau W, Schilling N, Shaffer PW, Chien WS (1978) Pathways of assimilation of [¹⁵N] N₂ and ¹⁵NH₄⁺ by cyanobacteria with and without heterocysts. *Journal of Bacteriology* **134**, 125–130.
- Mendez JM, Vega JM (1981) Purification and molecular properties of nitrite reductase from *Anabaena* sp. 7119. *Physiologia Plantarum* **52**, 7–14.
- Merida A, Leurentop L, Candau P, Florencio FJ (1990) Purification and properties of glutamine synthetases from the cyanobacteria *Synechocystis* sp. strain PCC 6803 and *Calothrix* sp. strain PCC 7601. *Journal of Bacteriology* **172**, 4732–4735.
- Mikami B, Ida S (1984) Purification and properties of ferredoxin nitrate reductase from the cyanobacterium *Plectonema boryanum*. *Biochimica et Biophysica Acta* **791**, 294–304.
- Moffett JW, Brand LE, Croot PL, Barbeau KA (1997) Cu speciation and cyanobacterial distribution in harbors subject to anthropogenic Cu inputs. *Limnology and Oceanography* **42**, 789–799.
- Morel FMM (2008) The co-evolution of phytoplankton and trace element cycles in the oceans. *Geobiology* **6**, 318–324.
- Morel FMM, Price NM (2003) The biogeochemical cycles of trace metals in the oceans. *Science* **300**, 944–947.
- Morel FMM, Milligan AJ, Saito MA (2003) Marine bioinorganic chemistry: the role of trace metals in the oceanic cycles of major nutrients. In *Treatise on Geochemistry* (eds Holland HD, Turekian KK). Elsevier Academic Press, London, UK, pp. 113–143.
- Moreno-Vivian C, Cabello P, Martinez-Luque M, Blasco R, Castillo F (1999) Prokaryotic nitrate reduction: molecular properties and functional distinction among bacterial nitrate reductases. *Journal of Bacteriology* **181**, 6573–6584.
- Muro-Pastor MI, Reyes JC, Florencio FJ (2005) Ammonium assimilation in cyanobacteria. *Photosynthesis Research* **83**, 135–150.
- Navarro F, Martin-Figueroa E, Candau P, Florencio FJ (2000) Ferredoxin-dependent iron-sulfur flavoprotein glutamate synthase (GlsF) from the cyanobacterium *Synechocystis* sp PCC 6803: Expression and assembly in *Escherichia coli*. *Archives of Biochemistry and Biophysics* **379**, 267–276.
- Navarro-González R, Molina MJ, Molina LT (1998) Nitrogen fixation by volcanic lightning in the early Earth. *Geophysical Research Letters* **25**, 3123–3126.
- Navarro-Gonzalez R, McKay CP, Mvondo DN (2001) A possible nitrogen crisis for Archaean life due to reduced nitrogen fixation by lightning. *Nature* **412**, 61–64.

- Nicholas DJD, Nason A (1954) Molybdenum and nitrate reductase II. Molybdenum as a constituent of nitrate reductase. *Journal of Biological Chemistry* **207**, 353–360.
- Nicholas DJD, Nason A, McElroy WD (1954) Molybdenum and nitrate reductase I. Effect of molybdenum deficiency on the *Neurospora* enzyme. *Journal of Biological Chemistry* **207**, 341–352.
- Normand P, Bousquet J (1989) Phylogeny of nitrogenase sequences in *Frankia* and other nitrogen-fixing microorganisms. *Journal of Molecular Evolution* **29**, 436–447.
- Oda Y, Samanta SK, Rey FE, Wu L, Liu X, Yan T, Zhou J, Harwood CS (2005) Functional genomic analysis of three nitrogenase isozymes in the photosynthetic bacterium *Rhodospseudomonas palustris*. *Journal of Bacteriology* **187**, 7784–7794.
- Ohkouchi N, Kashiyama Y, Kuroda J, Ogawa NO, Kitazato H (2006) An importance of diazotrophic cyanobacteria as a primary producer during Cretaceous Oceanic Anoxic Event 2. *Biogeosciences* **3**, 467–478.
- Oliveira L, Antia NJ (1984) Evidence of nickel ion requirement for autotrophic growth of a marine diatom with urea serving as nitrogen source. *British Phycological Journal* **19**, 125–134.
- Oliveira L, Antia NJ (1986a) Nickel ion requirements for autotrophic growth of several marine microalgae with urea serving as nitrogen source. *Canadian Journal of Fisheries and Aquatic Sciences* **43**, 2427–2433.
- Oliveira L, Antia NJ (1986b) Some observations on the urea-degrading enzyme of the diatom *Cyclotella cryptica* and the role of nickel in its production. *Journal of Plankton Research* **8**, 235–242.
- Orr J, Keefer LM, Keim P, Nguyen TD, Wellems T, Heinrikson RL, Haselkorn R (1981) Purification, physical characterization, and NH₂-terminal sequence of glutamine synthetase from the cyanobacterium *Anabaena* 7120. *Journal of Biological Chemistry* **256**, 13091–13098.
- Palenik B, Morel FMM (1990a) Amino acid utilization by marine phytoplankton – a novel mechanism. *Limnology and Oceanography* **35**, 260–269.
- Palenik B, Morel FMM (1990b) Comparison of cell-surface l-amino acid oxidases from several marine phytoplankton. *Marine Ecology Progress Series* **59**, 195–201.
- Palenik B, Morel FMM (1991) Amine oxidases of marine phytoplankton. *Applied and Environmental Microbiology* **57**, 2440–2443.
- Palinska KA, Jahns T, Rippka R, de Marsac NT (2000) *Prochlorococcus marinus* strain PCC 9511, a picoplanktonic cyanobacterium, synthesizes the smallest urease. *Microbiology* **146**, 3099–3107.
- Papineau D, Mojzsis SJ, Karhu JA, Marty B (2005) Nitrogen isotopic composition of ammoniated phyllosilicates: case studies from Precambrian metamorphosed sedimentary rocks. *Chemical Geology* **216**, 37–58.
- Pau RN, Mitchenall LA, Robson RL (1989) Genetic evidence for an *Azotobacter vinelandii* nitrogenase lacking molybdenum and vanadium. *Journal of Bacteriology* **171**, 124–129.
- Pearce CR, Cohen AS, Coe AL, Burton KW (2008) Molybdenum isotope evidence for global ocean anoxia coupled with perturbations to the carbon cycle during the Early Jurassic. *Geology* **36**, 231.
- Petri R, Imhoff JF (2000) The relationship of nitrate reducing bacteria on the basis of *narH* gene sequences and comparison of *narH* and 16S rDNA based phylogeny. *Systematic and Applied Microbiology* **23**, 47–57.
- Philippot L (2002) Denitrifying genes in bacterial and archaeal genomes. *Biochimica et Biophysica Acta-Gene Structure et Expression* **1577**, 355–376.
- Poulton SW, Fralick PW, Canfield DE (2004) The transition to a sulphidic ocean approximately 1.84 billion years ago. *Nature* **431**, 173–177.
- Price NM, Morel FMM (1991) Colimitation of phytoplankton growth by nickel and nitrogen. *Limnology and Oceanography* **36**, 1071–1077.
- Price NM, Andersen LF, Morel FMM (1991) Iron and nitrogen nutrition of equatorial Pacific plankton. *Deep-Sea Research Part A-Oceanographic Research Papers* **38**, 1361–1378.
- Quigg A, Finkel ZV, Irwin AJ, Rosenthal Y, Ho T-Y, Reinfelder JR, Schofield O, Morel FMM, Falkowski PG (2003) The evolutionary inheritance of elemental stoichiometry in marine phytoplankton. *Nature* **425**, 291–294.
- Rai AK (1989) Purification and properties of urease from a cyanobacterium *Anabaena doliolum*. *FEMS Microbiology Letters* **61**, 319–322.
- Rasmussen B, Fletcher IR, Brocks JJ, Kilburn MR (2008) Reassessing the first appearance of eukaryotes and cyanobacteria. *Nature* **455**, 1101–1104.
- Ravasio S, Dossena L, Martin-Figueroa E, Florencio FJ, Mattevi A, Morandi P, Curti B, Vanoni MA (2002) Properties of the recombinant ferredoxin-dependent glutamate synthase of *Synechocystis* PCC6803. Comparison with the *Azospirillum brasilense* NADPH-dependent enzyme and its isolated alpha subunit. *Biochemistry* **41**, 8120–8133.
- Raymond J (2005) The evolution of biological carbon and nitrogen cycling – a genomic perspective. *Reviews in Mineralogy and Geochemistry* **59**, 211–231.
- Raymond J, Segre D (2006) The effect of oxygen on biochemical networks and the evolution of complex life. *Science* **311**, 1764–1767.
- Raymond J, Siefert JL, Staples CR, Blankenship RE (2004) The natural history of nitrogen fixation. *Molecular Biology and Evolution* **21**, 541–554.
- Reade JPH, Dougherty LJ, Rogers LJ, Gallon JR (1999) Synthesis and proteolytic degradation of nitrogenase in cultures of the unicellular cyanobacterium *Gloeotheca* strain ATCC 27152. *Microbiology* **145**, 1749–1758.
- Rees TAV, Bekheet IA (1982) The role of nickel in urea assimilation by algae. *Planta* **156**, 385–387.
- Revill AT, Volkman JK, O'Leary T, Summons RE, Boreham CJ, Banks MR, Denwer K (1994) Hydrocarbon biomarkers, thermal maturity, and depositional setting of tasmanite oil shales from Tasmania, Australia. *Geochimica et Cosmochimica Acta* **58**, 3803–3822.
- Robson RL, Eady RR, Richardson TH, Miller RW, Hawkins M, Postgate JR (1986) The alternative nitrogenase of *Azotobacter chroococcum* is a vanadium enzyme. *Nature* **322**, 388–390.
- Romero LC, Galvan F, Vega JM (1987) Purification and properties of the siroheme-containing ferredoxin-nitrite reductase from *Chlamydomonas reinhardtii*. *Biochimica et Biophysica Acta* **914**, 55–63.
- de la Rosa MA (1983) Assimilatory nitrate reductase from the green alga *Ankistrodesmus braunii*. *Molecular and Cellular Biochemistry* **50**, 65–74.
- de la Rosa MA, Diez J, Vega JM, Losada M (1980) Purification and properties of assimilatory nitrate reductase [NAD(P)H] from *Ankistrodesmus braunii*. *FEBS Journal* **106**, 249–256.
- de la Rosa MA, Vega JM, Zumft WG (1981) Composition and structure of assimilatory nitrate reductase from *Ankistrodesmus braunii*. *Journal of Biological Chemistry* **256**, 5814–5819.
- Rouxel OJ, Bekker A, Edwards KJ (2005) Iron isotope constraints on the Archean and Paleoproterozoic ocean redox state. *Science* **307**, 1088–1091.

- Rubio LM, Ludden PW (2005) Maturation of nitrogenase: a biochemical puzzle. *Journal of Bacteriology* **187**, 405–414.
- Rubio LM, Ludden PW (2008) Biosynthesis of the iron-molybdenum cofactor of nitrogenase. *Annual Review of Microbiology* **62**, 93–111.
- Rubio LM, Flores E, Herrero A (1999) Molybdopterin guanine dinucleotide cofactor in *Synechococcus* sp. nitrate reductase: identification of *mobA* and isolation of a putative *moeB* gene. *FEBS Letters* **462**, 358–362.
- Rubio LM, Flores E, Herrero A (2002) Purification, cofactor analysis, and site-directed mutagenesis of *Synechococcus* ferredoxin-nitrate reductase. *Photosynthesis Research* **72**, 13–26.
- Saito MA, Sigman DM, Morel FMM (2003) The bioinorganic chemistry of the ancient ocean: the co-evolution of cyanobacterial metal requirements and biogeochemical cycles at the Archean-Proterozoic boundary? *Inorganica Chimica Acta* **356**, 308–318.
- Saito MA, Bertrand EM, Bulygin V, Moran D, Waterbury JB (2008) Proteomic analysis of the marine cyanobacterium *Synechococcus* WH8102 and implications for estimates of the cellular iron content. *Eos Trans. AGU* **89**(53), Fall Meet. Suppl., Abstract B31F–0358.
- Saltzman MR (2005) Phosphorus, nitrogen, and the redox evolution of the Paleozoic oceans. *Geology* **33**, 573–576.
- Sampaio M, Rowell P, Stewart WDP (1979) Purification and some properties of glutamine synthetase from the nitrogen-fixing cyanobacteria *Anabaena cylindrica* and a *Nostoc* sp. *Journal of General Microbiology* **111**, 181–191.
- Schemberg J, Schneider K, Demmer U, Warkentin E, Muller A, Ermler U (2007) Towards biological supramolecular chemistry: a variety of pocket-templated, individual metal oxide cluster nucleations in the cavity of a Mo/W-storage protein. *Angewandte Chemie International Edition* **119**, 2408–2413.
- Schneider K, Gollan U, Drottboom M, Selsemeier-Voigt S, Muller A (1997) Comparative biochemical characterization of the iron-only nitrogenase and the molybdenum nitrogenase from *Rhodobacter capsulatus*. *European Journal of Biochemistry* **244**, 789–800.
- Schoonen MAA, Xu Y (2001) Nitrogen reduction under hydrothermal vent conditions: Implications for the prebiotic synthesis of CHON compounds. *Astrobiology* **1**, 133–142.
- Schrautemeier B, Bohme H (1985) A distinct ferredoxin for nitrogen fixation isolated from heterocysts of the cyanobacterium *Anabaena variabilis*. *FEBS Letters* **184**, 304–308.
- Scott C, Lyons TW, Bekker A, Shen Y, Poulton SW, Chu X, Anbar AD (2008) Tracing the stepwise oxygenation of the Proterozoic ocean. *Nature* **452**, 456–459.
- Sharp JH (1983) The distribution of inorganic and organic nitrogen and dissolved and particulate nitrogen in the sea. In: *Nitrogen in the Marine Environment* (eds Carpenter EJ, Capone DG). Academic Press, New York, pp. 1–33.
- Shen Y, Buick R, Canfield DE (2001) Isotopic evidence for microbial sulphate reduction in the early Archaean era. *Nature* **410**, 77–81.
- Shen Y, Knoll AH, Walter MR (2003) Evidence for low sulphate and anoxia in a mid-Proterozoic marine basin. *Nature* **423**, 632–635.
- Shen Y, Zhang T, Hoffman PF (2008) On the coevolution of Ediacaran oceans and animals. *Proceedings of the National Academy of Sciences, USA* **105**, 7376.
- da Silva JJRF, Williams RJP (1991) *The Biological Chemistry of the Elements: the Inorganic Chemistry of Life*. Clarendon Press Oxford.
- Silver WS, Postgate JR (1973) Evolution of symbiotic nitrogen fixation. *Journal of Theoretical Biology* **40**, 1–10.
- Skipper L, Campbell WH, Mertens JA, Lowe DJ (2001) Pre-steady-state kinetic analysis of recombinant *Arabidopsis* NADH: nitrate reductase: rate-limiting processes in catalysis. *Journal of Biological Chemistry* **276**, 26995–27002.
- Smil V (2004) *Enriching the Earth: Fritz Haber, Carl Bosch, and the Transformation of World Food Production*, MIT Press, Cambridge, MA.
- Solomonson LP (1975) Purification of NADH-nitrate reductase by affinity chromatography. *Plant Physiology* **56**, 853–855.
- Solomonson LP, Lorimer GH, Hall RL, Borchers R, Bailey JL (1975) Reduced nicotinamide adenine dinucleotide-nitrate reductase of *Chlorella vulgaris*. Purification, prosthetic groups, and molecular properties. *Journal of Biological Chemistry* **250**, 4120–4127.
- Solomonson LP, Barber MJ, Robbins AP, Oaks A (1986) Functional domains of assimilatory NADH: nitrate reductase from *Chlorella*. *Journal of Biological Chemistry* **261**, 11290–11294.
- Stacey G, Tabita FR, Vanbaalen C (1977) Nitrogen and ammonia assimilation in cyanobacteria – purification of glutamine synthetase from *Anabaena* sp. strain CA. *Journal of Bacteriology* **132**, 596–603.
- Stal LJ, Zehr JP (2008) Cyanobacterial nitrogen fixation in the ocean: diversity, regulation, and ecology. In *The Cyanobacteria: Molecular Biology, Genomics and Evolution* (eds Herrero A, Flores E). Horizon Scientific Press, Norwich, UK, pp. 423–471.
- Steinberg RA (1937) Role of molybdenum in the utilization of ammonium and nitrate nitrogen by *Aspergillus niger*. *Journal of Agricultural Research* **55**, 891.
- Stolz JF, Basu P (2002) Evolution of nitrate reductase: molecular and structural variations on a common function. *ChemBiochemistry* **3**, 198–206.
- Sugawara K, Okabe S, Tanaka M (1961) Geochemistry of molybdenum in natural waters (II). *Journal of Earth Sciences* **9**, 114–128.
- Sumar N, Casselton PJ, McNally SF, Stewart GR (1984) Occurrence of isoenzymes of glutamine synthetase in the alga *Chlorella kessleri*. *Plant Physiology* **74**, 204–207.
- Summers DP, Chang S (1993) Prebiotic ammonia from reduction of nitrite by iron(II) on the early Earth. *Nature* **365**, 630–633.
- Suzuki A, Knaff DB (2005) Glutamate synthase: structural, mechanistic and regulatory properties, and role in the amino acid metabolism. *Photosynthesis Research* **83**, 191–217.
- Swamy U, Wang M, Tripathy JN, Kim SK, Hirasawa M, Knaff DB, Allen JP (2005) Structure of spinach nitrite reductase: implications for multi-electron reactions by the iron-sulfur: siroheme cofactor. *Biochemistry* **44**, 16054–16063.
- Syrett PJ, Alhouthy FAA (1984) The phylogenetic significance of the occurrence of urease urea amidolyase and glycollate oxidase glycollate dehydrogenase in green algae. *British Phycological Journal* **19**, 11–21.
- Tamagnini P, Axelsson R, Lindberg P, Oxelfelt F, Wunschiers R, Lindblad P (2002) Hydrogenases and hydrogen metabolism of cyanobacteria. *Microbiology and Molecular Biology Reviews* **66**, 1–20.
- Temple SJ, Vance CP, Stephen Gantt J (1998) Glutamate synthase and nitrogen assimilation. *Trends in Plant Science* **3**, 51–56.
- Thaivanich S, Incharoensakdi A (2007) Purification and characterization of nitrate reductase from the halotolerant cyanobacterium *Aphanothece halophytica*. *World Journal of Microbiology and Biotechnology* **23**, 85–92.
- Thiel T (1993) Characterization of genes for an alternative nitrogenase in the cyanobacterium *Anabaena variabilis*. *Journal of Bacteriology* **175**, 6276–6286.
- Thiel T, Pratte B, Zahalak M (2002) Transport of molybdate in the cyanobacterium *Anabaena variabilis* ATCC 29413. *Archives of Microbiology* **179**, 50–56.
- Tomitani A, Knoll AH, Cavanaugh CM, Ohno T (2006) The evolutionary diversification of cyanobacteria: molecular-phylogenetic and paleontological perspectives. *Proceedings of the National Academy of Sciences, USA* **103**, 5442–5447.

- Tuit C, Waterbury J, Ravizzaz G (2004) Diel variation of molybdenum and iron in marine diazotrophic cyanobacteria. *Limnology and Oceanography* **49**, 978–990.
- Valentin K, Kostrzewa M, Zetsche K (1993) Glutamate synthase is plastid-encoded in a red alga: implications for the evolution of glutamate synthases. *Plant Molecular Biology* **23**, 77–85.
- Vanoni MA, Curti B (1999) Glutamate synthase: a complex iron-sulfur flavoprotein. *Cellular and Molecular Life Sciences* **55**, 617–638.
- Vanoni MA, Dossena L, Heuvel RHH, Curti B (2005) Structure–function studies on the complex iron–sulfur flavoprotein glutamate synthase: the key enzyme of ammonia assimilation. *Photosynthesis Research* **83**, 219–238.
- Vega JMA, Herrera J, Aparicio PJ, Paneque A, Losada M (1971) Role of molybdenum in nitrate reduction by *Chlorella*. *Plant Physiology* **48**, 294–299.
- Walker J (1953) Inorganic micronutrient requirements of *Chlorella*. I. Requirements for calcium (or strontium), copper, and molybdenum. *Archives of Biochemistry and Biophysics* **46**, 1–11.
- Wallen DG, Cartier LD (1975) Molybdenum dependence, nitrate uptake and photosynthesis of freshwater plankton algae. *Journal of Phycology* **11**, 345–349.
- Wang QF, Li H, Post AF (2000) Nitrate assimilation genes of the marine diazotrophic, filamentous cyanobacterium *Trichodesmium* sp. strain WH9601. *Journal of Bacteriology* **182**, 1764–1767.
- Wille M, Nagler TF, Lehmann B, Schroder S, Kramers JD (2008) Hydrogen sulphide release to surface waters at the Precambrian/Cambrian boundary. *Nature* **453**, 767–769.
- Wolk CP, Thomas J, Shaffer PW, Austin SM, Galonsky A (1976) Pathway of nitrogen metabolism after fixation of ¹⁵N-labeled nitrogen gas by the cyanobacterium, *Anabaena cylindrica*. *Journal of Biological Chemistry* **251**, 5027–5034.
- Xie S, Pancost RD, Yin H, Wang H, Evershed RP (2005) Two episodes of microbial change coupled with Permo/Triassic faunal mass extinction. *Nature* **434**, 494–497.
- Xue HB, Kistler D, Sigg L (1995) Competition of copper and zinc for strong ligands in a eutrophic lake. *Limnology and Oceanography* **40**, 1142–1152.
- Xue HB, Jansen S, Prasch A, Sigg L (2001) Nickel speciation and complexation kinetics in freshwater by ligand exchange and DPCSV. *Environmental Science and Technology* **35**, 539–546.
- Yamaguchi KE (2002) Geochemistry of Archean–Paleoproterozoic black shales: early evolution of the atmosphere, oceans, and biosphere. Ph.D. Thesis. Pennsylvania State University, University Park, p. 456.
- Yuan HF, Wang CM, Kung HW (2001) Purification and characterization of glutamine synthetase from the unicellular cyanobacterium *Synechococcus* RF-1. *Botanical Bulletin of Academia Sinica* **42**, 23–33.
- Yung YL, McElroy MB (1979) Fixation of nitrogen in the prebiotic atmosphere. *Science* **203**, 1002–1004.
- Zahalak M, Pratte B, Werth KJ, Thiel T (2004) Molybdate transport and its effect on nitrogen utilization in the cyanobacterium *Anabaena variabilis* ATCC 29413. *Molecular Microbiology* **51**, 539–549.
- Zehr JP, Harris D, Dominic B, Salerno J (1997) Structural analysis of the *Trichodesmium* nitrogenase iron protein: implications for aerobic nitrogen fixation activity. *FEMS Microbiology Letters* **153**, 303–309.
- Zerle AL, House CH, Brantley SL (2005) Biogeochemical signatures through time as inferred from whole microbial genomes. *American Journal of Science* **305**, 467.
- Zerle AL, House CH, Cox RP, Canfield DE (2006) Metal limitation of cyanobacterial N₂ fixation and implications for the Precambrian nitrogen cycle. *Geobiology* **4**, 285–297.
- Zumft WG, Paneque A, Aparicio PJ, Losada M (1969) Mechanism of nitrate reduction in *Chlorella*. *Biochemical and Biophysical Research Communications* **36**, 980–986.
- Zumft WG, Aparicio PJ, Paneque A, Losada M (1970) Structural and functional role of FAD in the NADH-nitrate reducing system from *Chlorella*. *FEBS Letters* **9**, 157–160.

SUPPORTING INFORMATION

Additional Supporting Information may be found in the online version of this article:

Supplementary Text Figure 2. Detail of the construction of Figure 2 and of model organism species names.

Supplementary Text Figure 6. Details of the growth conditions of the experiment shown in Figure 6.

Please note: Wiley-Blackwell are not responsible for the content or functionality of any supporting materials supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.